

Extended Propagation of Powerful Laser Pulses in Focusing Kerr Media

V. M. Malkin and N. J. Fisch

Department of Astrophysical Sciences, Princeton University, Princeton, New Jersey 08540, USA

(Received 4 December 2015; published 20 September 2016)

Powerful incoherent laser pulses can propagate in focusing Kerr media much longer distances than can coherent pulses, due to the fast phase mixing that prevents transverse filamentation. This distance is limited by 4-wave scattering, which accumulates waves at small transverse wave numbers, where phase mixing is too slow to retain the incoherence and thus prevent the filamentation. However, we identify how this theoretical limit can be overcome by countering this accumulation through transverse heating of the pulse by random fluctuations of the refractive index. Thus, the laser pulse propagation distances are significantly extended, making feasible, in particular, the generation of unprecedentedly intense and powerful short laser pulses in a plasma by means of backward Raman amplification in new random laser regimes.

DOI: 10.1103/PhysRevLett.117.133901

The propagation of powerful laser pulses in focusing nonlinear media depends on the competition between transverse dispersive spreading and focusing nonlinearity [1–6]. For negligible nonlinearity, the dispersion would double the cross section of a coherent pulse of transverse size L_{\perp} within the Rayleigh length $L_R \sim L_{\perp}^2/\lambda$, where λ is the laser wavelength. A strong enough nonlinearity would noticeably reduce the pulse cross section, or even cause transverse filamentation, within the length $L_{SF} \sim v_g/\omega_{nl}$, where ω_{nl} is the nonlinear frequency shift and v_g is the group velocity of the pulse. For $L_R < L_{SF}$, the dispersive spreading outruns the self-focusing and suppresses it. We will consider focusing Kerr-like media for which the nonlinear frequency shift is proportional to the pulse intensity I , namely, $\omega_{nl} = -\alpha I$ with the positive coefficient $\alpha > 0$. Then, $L_R/L_{SF} \sim P_{ch}/P_{cr}$, where $P_{ch} \sim IL_{\perp}^2$ is the coherent pulse power and $P_{cr} \sim \lambda v_g/\alpha$ is the critical power of the self-focusing.

A coherent laser pulse of power P much greater than the critical power, $P \gg P_{cr}$, would experience transverse filamentation within a propagation length not much exceeding the self-focusing length L_{SF} . In contrast to this, incoherent laser pulses could traverse focusing Kerr-like media, remaining statistically uniform in the transverse directions, at arbitrarily large powers. What actually matters for the nonfilamentation of incoherent pulses is not the total power P , but the coherent subpower $P_{ch} \sim IL_{\perp}^2$ located within the transverse correlation length L_{\perp} . If this subpower is much smaller than the critical power, $P_{ch} \ll P_{cr}$, then the Rayleigh length is much shorter than the self-focusing length, so that nonlinear phase shift accumulated within the Rayleigh length is small, $\phi_{nl} \sim \omega_{nl}L_R/v_g \sim P_{ch}/P_{cr} \ll 1$.

The small parameter $P_{ch}/P_{cr} \ll 1$ enables a kind of perturbation theory. Physically, this small parameter implies that the phase mixing of different waves occurs much faster than the nonlinear interaction. In the zero-order

approximation, the phases of the different waves are random. Statistical averaging over nearly random phases leads to a closed evolution equation for the pair correlation function of the waves. This equation is usually referred to as a kinetic equation for the waves. The pair correlation function of the waves in the Wigner representation is usually referred to as the wave spectral density. In general, the kinetic equation for the waves contains terms linear, quadratic, cubic, etc. in the wave spectral density.

The linear term describes wave propagation with group velocities and scattering on inhomogeneities of the medium. For a pulse with negligible spread of group velocities in a conservative uniform medium, the wave spectral density does not change in the reference frame moving with the group velocity of the pulse.

The quadratic term in the kinetic equation for the waves gives a linear contribution to the wave spectral density variation rate. This contribution comes from the statistically averaged nonlinear frequency shift, proportional to the pulse intensity. The corresponding nonlinear phase shift can bend phase fronts and cause the self-focusing of incoherent pulses in a way similar to that of coherent pulses. The evolution of the transverse size (radius) of an incoherent laser pulse within such a model was discussed in many papers (see, for instance, Refs. [7,8]). This approximation is sufficient for Kerr media in which the nonlinear frequency shift is proportional to the laser field intensity integrated over time or space, because such an integration effectively accomplishes statistical averaging (see, for instance, Ref. [9]). However, for the Kerr media of interest here in which the nonlinear frequency shift is proportional to the laser field intensity itself, higher order nonlinear processes can be important.

In fact, the higher order nonlinear processes appear to be dominant for sufficiently powerful incoherent laser pulses statistically nearly uniform in the transverse directions over many correlation lengths L_{\perp} . For statistically uniform

pulses, the averaged nonlinear frequency shift does not vary in the transverse directions, so that there is no bending of the phase fronts, and the quadratic term in the kinetic equation for the waves is zero. The dominant nonlinear term in the kinetic equation for the waves is then the cubic term.

More accurately, the cubic term in the kinetic equation for the waves, which gives the quadratic contribution to the wave spectral density variation rate, contains the small factor $P_{\text{ch}}^2/P_{\text{cr}}^2$. The quadratic term in the kinetic equation for the waves, which gives the linear contribution to the wave spectral density variation rate, contains the small factor $f_I P_{\text{ch}}/P_{\text{cr}}$, where f_I is the small relative variation of the wave spectral density within the transverse correlation length L_{\perp} . For a smooth wave spectral density nonmodulated in the transverse directions, the typical f_I is about the ratio of L_{\perp} to the pulse aperture, $f_I \sim \sqrt{P_{\text{ch}}/P}$. The cubic process is much faster than the quadratic process for $P \gg P_{\text{cr}}^2/P_{\text{ch}}$. Of interest here are pulses of very high powers P , perhaps exceeding the critical power P_{cr} by a factor of 10^6 , while, to secure the pulse incoherence, the ratio $P_{\text{cr}}/P_{\text{ch}}$ needs to be just somewhat larger than 1. The effect of the quadratic term in the kinetic equation for the waves on the propagation of such powerful pulses is then negligible.

The cubic term in the kinetic equation for the waves is associated with the 4-wave scattering. The evolution of the wave spectral density due to the 4-wave scattering was the subject of a longstanding theoretical controversy, as recounted and resolved analytically in Ref. [10]. That analytical resolution was recently supported numerically [11]. Being the same for focusing and defocusing Kerr nonlinearities, the 4-wave kinetic equation cannot produce self-focusing effects directly. However, it tends to accumulate waves at small transverse wave numbers, where the phase mixing is too slow to retain the incoherence. Therefore, the input pulse randomization (which might be arranged, say, by techniques of the type in Refs. [12–14]) is not sufficient to achieve pulse propagation over distances much exceeding the 4-wave scattering length.

Here, we propose to use an ongoing randomization to prevent wave accumulation at small transverse wave numbers, thus extending the propagation of powerful laser pulses in focusing Kerr media much beyond the known theoretical limit [10]. The ongoing randomization can be accomplished through natural ray diffusion in media with random fluctuations of the refractive index. The substantially extended propagation lengths could make feasible a new class of random lasers [15–17], based on backward Raman amplification [18–22] and capable of reaching relativistic nonfocused intensities in plasmas. This would exploit the random density fluctuations inherently present in plasmas and hitherto considered harmful to backward Raman amplification.

Since the wave spectral density variation rate, associated with the 4-wave scattering, contains the small factor

$P_{\text{ch}}^2/P_{\text{cr}}^2$ compared to the rate of phase mixing, the inverse length of 4-wave scattering is

$$L_{4w}^{-1} \sim L_R^{-1} P_{\text{ch}}^2/P_{\text{cr}}^2 \sim L_R/L_{\text{SF}}^2. \quad (1)$$

The 4-wave scattering of two waves with transverse wave numbers $k_{\perp} \sim 2\pi/L_{\perp}$ into two new waves typically produces a wave of larger transverse wave number along with a wave of smaller transverse wave number. Further 4-wave scattering produces waves of even larger transverse wave numbers, so that the wave distribution spreads over a circle of growing radius $k_{\perp M}$ in the k_{\perp} plane. At the same time, the number of waves with small k_{\perp} increases. For smaller k_{\perp} , the 4-wave scattering is faster $L_{4w}^{-1} \propto L_R \propto L_{\perp}^2 \propto k_{\perp}^{-2}$, and it tends to establish the wave spectral density close to the equilibrium Rayleigh-Jeans distribution inside the circle $k_{\perp} \lesssim k_{\perp M}$:

$$N_{k_{\perp}} \approx \frac{T(\Lambda_{k_{\perp}})}{k_{\perp}^2 + k_{\perp m}^2}, \quad \Lambda_{k_{\perp}} = \ln \left(1 + \frac{k_{\perp}^2}{k_{\perp m}^2} \right). \quad (2)$$

Here, $k_{\perp m}^2$ is the “chemical potential” and $T(\Lambda_{k_{\perp}})$ is the “local temperature,” which can logarithmically slowly depend on k_{\perp} to provide nonzero fluxes across the spectrum. This multiscale distribution has roughly $\Lambda_{k_{\perp M}}$ different populated scales, each of which carries just a small fraction of total pulse intensity I . Such an optical turbulence does not satisfy the classical Kolmogorov hypothesis of spectral locality of interactions. There is, however, a more general kind of locality [10] that enables expressing k_{\perp} integrals in the 4-wave kinetic equation by explicit formulas local in k_{\perp} , so that Eq. (1) is modified for the multiscale spectrum as follows:

$$L_{4wk_{\perp}}^{-1} \sim f'(\Lambda_{k_{\perp}})f(\Lambda_{k_{\perp}})L_{Rk_{\perp}}/L_{\text{SF}}^2, \\ f(\Lambda_{k_{\perp}}) = \int_{\xi \lesssim \Lambda_{k_{\perp}}} T(\xi) d\xi / \int_{\xi \lesssim \Lambda_{k_{\perp M}}} T(\xi) d\xi. \quad (3)$$

Here, $f(\Lambda_{k_{\perp}})$ is the fraction of the pulse intensity carried by waves with transverse wave numbers not exceeding k_{\perp} . A simplified explanation of the factor $f'f$ entering here is that, being quadratic in wave intensity, the rate of 4-wave scattering contains the product of two wave intensities; one of the contributing waves has a wave number about k_{\perp} , and thus carries the intensity $\sim f'(\Lambda_{k_{\perp}})I$, while the second wave may have any wave number smaller than k_{\perp} , and thus carries the intensity $\sim f(\Lambda_{k_{\perp}})I$. Therefore, the rate of 4-wave scattering is modified by the factor $\sim f'f$. The 4-wave scattering inverse length $L_{4wk_{\perp}}^{-1}$ relates to its initial value L_{40}^{-1} as

$$\frac{L_{40}}{L_{4wk_{\perp}}} \sim \frac{I^2 k_{\perp 0}^2}{I_0^2 k_{\perp}^2} f'(\Lambda_{k_{\perp}})f(\Lambda_{k_{\perp}}), \quad L_{40} \sim \frac{L_{\text{SF}0}^2}{L_{Rk_{\perp 0}}}. \quad (4)$$

The applicability condition of the random phase approximation, $L_{4wk_{\perp}}^{-1} < L_{Rk_{\perp}}^{-1}$, is most restrictive at $k_{\perp} \sim k_{\perp m}$.

There, $f(\Lambda_{k_{\perp m}}) \sim f'(\Lambda_{k_{\perp m}})$ and $L_{4wk_{\perp m}}^{-1} \sim f'^2(\Lambda_{k_{\perp m}})L_{Rk_{\perp m}}/L_{SF}^2$, so that the applicability condition reduces to $f'(\Lambda_{k_{\perp m}})L_{Rk_{\perp m}}/L_{SF} < 1$. It simply means that the Rayleigh length of the waves with $k_{\perp} \sim k_{\perp m}$ must be shorter than the self-focusing length of these waves themselves, regardless of the other parts of the spectrum. This can be rewritten in the form

$$f'(\Lambda_{k_{\perp m}})(k_{\perp 0}^2/k_{\perp m}^2)(I/I_0) < P_{cr}/P_{ch0}. \quad (5)$$

The growth of $k_{\perp M}^2$ is described by the equation

$$\frac{dk_{\perp M}^2}{dz} \sim \frac{k_{\perp M}^2}{L_{4wk_{\perp M}}} \sim \frac{I^2 k_{\perp 0}^2}{I_0^2 L_{40}} f'(\Lambda_{k_{\perp M}}). \quad (6)$$

As long as other effects do not cause fluxes, the 4-wave scattering establishes nearly the same local temperature across the spectrum, $T(\Lambda_{k_{\perp}}) \approx T(\Lambda_{k_{\perp M}}) = T$. It follows then that $f'(\Lambda_{k_{\perp}}) \approx 1/\Lambda_{k_{\perp M}}$ and $f(\Lambda_{k_{\perp}}) \approx \Lambda_{k_{\perp}}/\Lambda_{k_{\perp M}}$. The intensity I may change due to the longitudinal stretching or contraction, or pumping or damping, of the pulse. Even absent the group velocity dispersion and pumping or damping, the short pulse duration may be strongly affected by small nonparaxial corrections to the longitudinal component of the group velocity $v_{g\parallel}$. The relative reduction of $v_{g\parallel}$, $\delta v_{g\parallel}/v_{g\parallel} \approx k_{\perp}^2/(2k_{\parallel}^2)$, is largest for waves with $k_{\perp} \sim k_{\perp M}$. It may cause pulse stretching $\delta L \sim z k_{\perp M}^2/(2k_{\parallel}^2)$, which can be neglected only for pulses of sufficiently large length L ,

$$L > z k_{\perp M}^2/(2k_{\parallel}^2). \quad (7)$$

Consider, first, the regimes with $I \sim I_0$. Since $I = \int dk_{\perp} N_{k_{\perp}} \sim T \Lambda_{k_{\perp M}}$, it follows $T \sim I_0/\Lambda_{k_{\perp M}}$. The 4-wave scattering itself conserves also the ‘‘transverse energy’’ $\mathcal{E}_{\perp} = \int dk_{\perp} N_{k_{\perp}} k_{\perp}^2$. The major contribution to this integral comes from waves with $k_{\perp} \sim k_{\perp M}$, so that $\mathcal{E}_{\perp} \sim k_{\perp M}^2 T \sim I_0 k_{\perp M}^2/\Lambda_{k_{\perp M}}$. As long as \mathcal{E}_{\perp} is conserved, it follows that $\Lambda_{k_{\perp M}} \sim k_{\perp M}^2/k_{\perp 0}^2$. Equation (6) gives then $k_{\perp M}^4/k_{\perp 0}^4 \sim 2z/L_{40}$, so that

$$\ln(k_{\perp M}^2/k_{\perp 0}^2) \approx \Lambda_{k_{\perp M}} \sim k_{\perp M}^2/k_{\perp 0}^2 \sim (2z/L_{40})^{1/2}. \quad (8)$$

The applicability condition (5) takes now the form

$$\Lambda_{k_{\perp M}} < \Lambda_*, \quad \exp(\Lambda_*)/\Lambda_*^2 \sim P_{cr}/P_{ch0}. \quad (9)$$

It can be satisfied only up to z not much exceeding L_{40} , namely $z < z_* \sim L_{40}\Lambda_*^2/2$. To overcome this previous theoretical limit [10], the ray diffusion on the medium inhomogeneities should stop the exponential decrease of $k_{\perp m}$ before condition (9) is violated. The diffusion spreads the ray transverse wave vectors according to $(\delta k_{\perp})^2 = Dz$. It starts affecting the previous regime when (δk_{\perp}) matches $k_{\perp m}$. This occurs at $\Lambda_{k_{\perp M}} = \Lambda_D$, such that $\Lambda_D \exp(\Lambda_D) \sim 2k_{\perp 0}^2 L_{40}^{-1}/D$. The applicability condition (9) is satisfied for

$$D > D_* \sim 2k_{\perp 0}^2 L_{40}^{-1} \Lambda_*^{-1} \exp(-\Lambda_*). \quad (10)$$

For such D , a condensate does not form; rather, $k_{\perp m}^2$ passes the minimum Dz_D at $z_D \sim L_{40}\Lambda_D^2/2 < z_*$, and then grows at $z > z_D$ according to

$$k_{\perp m}^2 \sim Dz. \quad (11)$$

This increases the local temperature at $k_{\perp} \sim k_{\perp m}$, but the increase is quickly saturated by the faster 4-wave scattering, which produces a flux of waves towards larger k_{\perp} up to $k_{\perp M}$. As a result, a fixed profile of the local temperature $T(\Lambda_{k_{\perp}})$ sets up across the spectrum. The entire spectrum evolves then in a self-similar way, as a function of just $k_{\perp}/k_{\perp m} \propto k_{\perp}/\sqrt{z}$, rather than k_{\perp} and z separately. It implies $L_{4wk_{\perp}} \sim z \propto k_{\perp}^2$ across the spectrum, which, according to Eq. (4), means $f'(\Lambda_{k_{\perp}})f(\Lambda_{k_{\perp}}) = \text{const}$. The solution of this equation is

$$f(\Lambda_{k_{\perp}}) \approx \Lambda_{k_{\perp}}^{1/2} \Lambda_{k_{\perp M}}^{-1/2}. \quad (12)$$

Integration of Eq. (6) gives

$$k_{\perp M}^2/k_{\perp 0}^2 \sim \Lambda_{k_{\perp M}}^{-1} z/L_{40}, \quad \Lambda_{k_{\perp M}} \approx \Lambda_D. \quad (13)$$

This regime extends until the applicability condition (7) is violated and the longitudinal stretching of the pulse becomes important. At larger z , the pulse intensity decreases due to the longitudinal stretching, but the fluence $w = IL$ does not change, absent pumping or damping. The stretching is described by the equation

$$dL/dz \sim k_{\perp M}^2/(2k_{\parallel}^2). \quad (14)$$

Though the stretching is mainly due to the waves with $k_{\perp} \sim k_{\perp M}$ trailing, the 4-wave scattering quickly sets up nearly the same transverse spectrum at all cross sections within the pulse length. Therefore, formula (12) remains valid. The variation of $k_{\perp M}^2$ is described by Eq. (6). For a fixed fluence w , the relative variation of $k_{\perp M}^2$ is small, so that the stretching proceeds linearly $L \propto z$. For a fixed pumping, the fluence grows linearly, $w \propto z$, which gives $L \propto z^{4/3}$ and $k_{\perp M}^2 \propto z^{1/3}$. Waves with $k_{\perp} \sim k_{\perp M}$ are not directly affected by the ray diffusion, so that the spectrum remains dominated by the 4-wave scattering and multi-scale, as long as $k_{\perp M}^2 > Dz$. When this condition is violated, the 4-wave scattering becomes relatively small and the propagation regime becomes nearly linear.

Consider now the calculation of the ray diffusion coefficient D for a medium with statistically uniform random inhomogeneities of the refractive index having a typical relative amplitude of fluctuations f_r , a transverse correlation length $L_{\perp r}$, and a longitudinal correlation length $L_{\parallel r}$. Let the group and phase velocities of the laser pulse be comparable. Then, according to the Hamilton equations for the rays, the transverse wave number typically changes, during the ray passing a single fluctuation, by

$\delta k_{\perp} \sim k_{\parallel} f_r L_{\parallel r} / L_{\perp r}$, for not too small an aspect ratio $L_{\perp r} / L_{\parallel r} \gtrsim k_{\perp} / k_{\parallel}$. The respective diffusion coefficient is

$$D \sim \delta k_{\perp}^2 / L_{\parallel r} \sim k_{\parallel}^2 f_r^2 L_{\parallel r} / L_{\perp r}^2. \quad (15)$$

To prevent Bose-Einstein condensation, this diffusion coefficient should exceed the threshold D_* . The fluctuation amplitude needed is then

$$f_r > f_{r*} \sim \frac{L_{\perp r} k_{\perp 0} \sqrt{2} \exp(-\Lambda_*/2)}{k_{\parallel} \sqrt{L_{\parallel r} L_{40} \Lambda_*}}. \quad (16)$$

In particular, for a plasma with fluctuations of the electron concentration δn and a plasma-to-laser frequency ratio $f_e = \omega_e / \omega = \lambda / \lambda_e \ll 1$ (where $\lambda_e = 2\pi c / \omega_e$), the refractive index fluctuation amplitude is $f_r = f_e^2 \delta n / (2n_0)$. The fluctuations δn needed to prevent Bose-Einstein condensate formation is then

$$\frac{\delta n}{n_0} > \frac{\delta n_*}{n_0} \sim \frac{L_{\perp r} \lambda_e^2 (2\Lambda_*)^{3/2} \exp(-3\Lambda_*/2)}{(L_{\parallel r} \lambda)^{1/2} L_{\perp 0}^2}. \quad (17)$$

Introducing such fluctuations is key to extending the laser propagation in a plasma, making possible the amplification of short laser pulses to unprecedented fluences. Pulses of durations too short to noticeably move ions, or to heat the plasma, can be amplified to ultrahigh intensities, since the ponderomotive and thermal nonlinearities, leading to self-focusing and filamentation, do not develop. What limits the propagation and amplification of such short pulses in the plasma is the relativistic electron nonlinearity, which comes from the electron mass dependence of the electron quiver velocity, $m_e = m / \sqrt{1 - v^2/c^2} \approx m(1 + 0.5v^2/c^2)$. The normalized quiver energy, averaged over the laser period, $\overline{v^2}/c^2$ can be expressed in terms of the laser intensity I as $\overline{v^2}/c^2 = 4\pi e^2 I / (m^2 c^3 \omega^2)$. The laser frequency shift due to the relativistic electron nonlinearity is $\omega_{nl} \approx -\overline{v^2} \omega_e^2 / (4c^2 \omega) = -\alpha I$, $\alpha = \pi e^2 \omega_e^2 / (m^2 c^3 \omega^3) = e^2 \lambda^3 / (2m^2 c^4 \lambda_e^2)$. The critical self-focusing power is then

$$P_{cr} \sim \lambda v_g / \alpha \approx \mathcal{P} \lambda_e^2 / \lambda^2, \quad \mathcal{P} = 2m^2 c^5 / e^2 \approx 17 \text{ GW}, \quad (18)$$

which well agrees with the standard value of P_{cr} for axisymmetric laser pulses in a plasma [23–25].

The relativistic electron nonlinearity limits the classical π -pulse regime of backward Raman amplification in a plasma [20,21,26–29], during which the pulse amplitude grows proportionally to z and the pulse length contracts inversely. This ends when the self-focusing length, decreasing in the π -pulse regime like z^{-2} , matches z . Then, the leading spike growth saturates and the transverse filamentation instability becomes dangerous. High-quality coherent pulses may propagate somewhat further without the filamentation, but no more than a few times this distance,

until the filamentation instability makes enough exponentiations to grow from small initial imperfections of the wave fronts. Such a modestly extended propagation allows us to amplify coherent pulses to higher fluences if not intensities [28]. However, incoherent pulses can propagate much longer distances, remaining statistically uniform in the transverse directions. Since the wave fronts of incoherent pulses are not smooth at all, the filamentation instability must be suppressed even before the leading amplified spike saturates at the intensity [27]

$$I_0 \approx \frac{\mathcal{P}}{\lambda_e \lambda} \left(\frac{I_{p0}}{I_{br}} \right)^{2/3}, \quad I_{br} = \frac{\pi \mathcal{P} \lambda}{32 \lambda_e^3}. \quad (19)$$

Here, I_{p0} is the incident pump pulse intensity, and I_{br} is its value at which wave breaking occurs of the resonant Langmuir wave, mediating energy transfer from the pump to the amplified pulse; it is assumed $I_{br} > I_{p0}$, because the energy transfer efficiency significantly drops at $I_{br} \ll I_{p0}$ [20,26,30,31]. As seen from Eq. (19), the energy compression ratio is $I_0 / I_{p0} \approx 10 (I_{br} / I_{p0})^{1/3} \lambda_e^2 / \lambda^2 > 1000$, for $\lambda_e / \lambda > 10$. This is also the ratio of the consumed pump length $2z_0$ to the compressed pulse length $L_0 \sim \lambda_e$. Within the propagation length z_0 , the pulse longitudinal stretching due to the finite angular spread is $\delta L_0 \sim z_0 \lambda^2 / (2L_{\perp 0}^2)$. Since $z_0 \sim L_0 I_0 / I_{p0}$ and $L_{\perp 0}^2 \sim I_0 / P_{ch0}$, it follows that $\delta L_0 / L_0 \sim 5 P_{cr} \lambda / (2 P_{ch0} \lambda_e) (I_{p0} / I_{br})^{1/3}$. For $P_{ch0} / P_{cr} > 5 \lambda / (2 \lambda_e) (I_{p0} / I_{br})^{1/3}$, this stretching is small, so that the π -pulse stage is not much affected by the finite angular spread. Substitution of $L_{\perp 0}^2 \sim I_{p0} / P_{ch0} \sim \lambda \lambda_e^{-3} (I_{p0} / I_{br})^{2/3} P_{cr} / P_{ch0}$ into Eq. (17), and using Eq. (9) to express P_{cr} / P_{ch0} in terms of Λ_* gives

$$\frac{\delta n}{n_0} > \frac{\delta n_*}{n_0} \sim \frac{2^{3/2} \exp(-\Lambda_*/2) L_{\perp r} \lambda^{1/2}}{\Lambda_*^{1/2} L_{\parallel r}^{1/2} \lambda_e} \left(\frac{I_{p0}}{I_{br}} \right)^{2/3}. \quad (20)$$

If, for example, $L_{\parallel r} = L_{\perp r} = \lambda_e = 25\lambda$, and $I_{p0} \sim I_{br}$, then $\delta n_* / n_0 \sim 15\%$ for $\Lambda_* = 2$, $\delta n_* / n_0 \sim 4\%$ for $\Lambda_* = 4$, and $\delta n_* / n_0 \sim 1\%$ for $\Lambda_* = 6$. For $\lambda = 0.3 \mu\text{m}$, the intensity I_0 is $I_0 \approx 0.8 \times 10^{18} \text{ W/cm}^2$, and the plasma concentration is $n_0 = \pi m c^2 (e \lambda_e)^{-2} \approx 2 \times 10^{19} \text{ cm}^{-3}$. Such a plasma might be produced by the ionization of dense aerosols with the droplets as small as a few microns [32,33].

With the filamentation instability suppressed by the pulse incoherence, backward Raman amplification can proceed over much larger distances and produce pulses of much higher fluences, if not intensities, than thought earlier. The high-energy output can be converted into an unprecedentedly intense and well-focused coherent laser pulse by means of backward Raman amplification in a thin dense plasma layer, as described in the two-step scheme [22]. The first step output is used as the pump at the second step. The transverse filamentation instability of the coherent amplified pulse does not pose a serious problem at the

second step, because of the plasma's short length and high density.

In summary, the key findings here are as follows: (a) Powerful laser pulse scattering on random fluctuations of the refractive index suppresses the transverse filamentation instability and enables propagation distances significantly exceeding the previous theoretical limit in focusing Kerr media, and (b) This scattering may now be used to overcome the transverse filamentation instability, the hitherto key limitation to producing laser pulses of unprecedented high intensities and powers through backward Raman amplification in a plasma.

This work was supported by DTRA HDTRA1-11-1-0037, by NSF PHY-1202162, and by the NNSA SSAA Program under Grant No. DENA0002948.

-
- [1] R. Y. Chiao, E. Garmire, and C. H. Townes, Self-Trapping of Optical Beams, *Phys. Rev. Lett.* **13**, 479 (1964).
- [2] V. I. Talanov, On self-focusing of electromagnetic waves in nonlinear mediums, *Izvestiya Vuzov. Radiofizika* **7**, 564 (1964).
- [3] P. L. Kelley, Self-Focusing of Optical Beams, *Phys. Rev. Lett.* **15**, 1005 (1965).
- [4] V. I. Bespalov and V. I. Talanov, Filamentary structure of light beams in nonlinear liquids, *JETP Lett.* **3**, 307 (1966).
- [5] S. A. Akhmanov, A. P. Sukhorukov, and R. V. Khokhlov, Self focusing and diffraction of light in a nonlinear medium, *Sov. Phys. Usp.* **10**, 609 (1968).
- [6] V. M. Malkin, On the analytical theory for stationary self-focusing of radiation, *Physica D (Amsterdam)* **64**, 251 (1993).
- [7] G. A. Pasmanik, Self-interaction of incoherent light beams, *Zh. Eksp. Teor. Fiz.* **66**, 490 (1974) [*Sov. Phys. JETP* **39**, 234 (1974)].
- [8] V. A. Aleshkevich, G. D. Kozhoridze, and A. N. Matveev, Self-action of partly coherent laser radiation, *Sov. Phys. Usp.* **34**, 777 (1991).
- [9] O. Bang, D. Edmondson, and W. Królikowski, Collapse of Incoherent Light Beams in Inertial Bulk Kerr Media, *Phys. Rev. Lett.* **83**, 5479 (1999).
- [10] V. M. Malkin, Kolmogorov and Nonstationary Spectra of Optical Turbulence, *Phys. Rev. Lett.* **76**, 4524 (1996).
- [11] G. Falkovich and N. Vladimirova, Cascades in nonlocal turbulence, *Phys. Rev. E* **91**, 041201 (2015).
- [12] R. H. Lehmberg and S. P. Obenschain, Use of induced spatial incoherence for uniform illumination of laser fusion targets, *Opt. Commun.* **46**, 27 (1983).
- [13] Y. Kato, K. Mima, N. Miyanaga, S. Arinaga, Y. Kitagawa, M. Nakatsuka, and C. Yamanaka, Random phasing of high-power lasers for uniform target acceleration and plasma-instability suppression, *Phys. Rev. Lett.* **53**, 1057 (1984).
- [14] S. Skupsky, R. W. Short, T. Kessler, R. S. Craxton, S. Letzring, and J. M. Soures, Improved laser-beam uniformity using the angular dispersion of frequency-modulated light, *J. Appl. Phys.* **66**, 3456 (1989).
- [15] D. Wiersma, The physics and applications of random lasers, *Nat. Phys.* **4**, 359 (2008).
- [16] S. Turitsyn, S. Babin, A. El-Taher, P. Harper, D. Churkin, S. Kablukov, J. Ania-Castn, V. Karalekas, and E. Podivilov, Random distributed feedback fibre laser, *Nat. Photonics* **4**, 231 (2010).
- [17] D. Churkin, I. Kolokolov, E. Podivilov, I. Vatik, M. Nikulin, S. Vergeles, I. Terekhov, V. Lebedev, G. Falkovich, S. Babin, and S. Turitsyn, Wave kinetics of random fibre lasers, *Nat. Commun.* **2**, 6214 (2015).
- [18] M. Maier, W. Kaiser, and J. A. Giordmaine, Intense Light Bursts in the Stimulated Raman Effect, *Phys. Rev. Lett.* **17**, 1275 (1966).
- [19] M. Maier, W. Kaiser, and J. A. Giordmaine, Backward stimulated Raman scattering, *Phys. Rev.* **177**, 580 (1969).
- [20] V. M. Malkin, G. Shvets, and N. J. Fisch, Fast Compression of Laser Beams to Highly Overcritical Powers, *Phys. Rev. Lett.* **82**, 4448 (1999).
- [21] V. M. Malkin, G. Shvets, and N. J. Fisch, Ultra-powerful compact amplifiers for short laser pulses, *Phys. Plasmas* **7**, 2232 (2000).
- [22] V. M. Malkin and N. J. Fisch, Manipulating ultra-intense laser pulses in plasmas, *Phys. Plasmas* **12**, 044507 (2005).
- [23] A. G. Litvak, Finite-amplitude wave beams in a magnetoactive plasma, *Zh. Eksp. Teor. Fiz.* **57**, 629 (1969) [*Sov. Phys. JETP* **30**, 344 (1970)].
- [24] C. Max, J. Arons, and A. B. Langdon, Self-Modulation and Self-Focusing of Electromagnetic Waves in Plasmas, *Phys. Rev. Lett.* **33**, 209 (1974).
- [25] G.-Z. Sun, E. Ott, Y. C. Lee, and P. Guzdar, Self-focusing of short intense pulses in plasmas, *Phys. Fluids* **30**, 526 (1987).
- [26] V. M. Malkin and N. J. Fisch, Key plasma parameters for resonant backward Raman amplification in plasma, *Eur. Phys. J. Spec. Top.* **223**, 1157 (2014).
- [27] V. M. Malkin, Z. Toroker, and N. J. Fisch, Saturation of the leading spike growth in backward Raman amplifiers, *Phys. Plasmas* **21**, 093112 (2014).
- [28] V. M. Malkin, Z. Toroker, and N. J. Fisch, Exceeding the leading spike intensity and fluence limits in backward Raman amplifiers, *Phys. Rev. E* **90**, 063110 (2014).
- [29] I. Barth, Z. Toroker, A. A. Balakin, and N. J. Fisch, Beyond nonlinear saturation of backward Raman amplifiers, *Phys. Rev. E* **93**, 063210 (2016).
- [30] Z. Toroker, V. M. Malkin, and N. J. Fisch, Backward Raman amplification in the Langmuir wavebreaking regime, *Phys. Plasmas* **21**, 113110 (2014).
- [31] M. Edwards, Z. Toroker, J. Mikhailova, and N. Fisch, The efficiency of Raman amplification in the wavebreaking regime, *Phys. Plasmas* **22**, 074501 (2015).
- [32] M. J. Hay, E. J. Valeo, and N. J. Fisch, Geometrical Optics of Dense Aerosols: Forming Dense Plasma Slabs, *Phys. Rev. Lett.* **111**, 188301 (2013).
- [33] D. Ruiz, L. Gunderson, M. Hay, E. Merino, E. Valeo, S. Zweben, and N. Fisch, Aerodynamic focusing of high-density aerosols, *J. Aerosol Sci.* **76**, 115 (2014).