On the Saturated State of the Nonlinear Small-Scale Dynamo

A. A. Schekochihin,[1,*] S. C. Cowley,[1] S. F. Taylor,[1] G. W. Hammett,[2] J. L. Maron,[3,4] and J. C. McWilliams[5]

[1] *Plasma Physics Group, Imperial College, Blackett Laboratory, Prince Consort Road, London SW7 2BW, UK*
[2] *Plasma Physics Laboratory, Princeton University, Princeton, NJ 08543*
[3] *Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627*
[4] *Center for Magnetic Reconnection Studies, Department of Physics and Astronomy, University of Iowa, Iowa City, IA 52242*
[5] *Department of Atmospheric Sciences, UCLA, Los Angeles, CA 90095-1565*

(Dated: August 14, 2003)

We consider the problem of incompressible, forced, nonhelical, homogeneous and isotropic MHD turbulence with no mean magnetic field and large magnetic Prandtl number. This type of MHD turbulence is the end state of the turbulent dynamo, which generates folded fields with small-scale direction reversals. We propose a model in which saturation is achieved as a result of the velocity statistics becoming anisotropic with respect to the local direction of the folded fields. The model combines the effects of weakened stretching and quasi-two-dimensional mixing and produces folded-field spectra that are in remarkable agreement with numerical results at least in the case of a one-scale flow. We conjecture that the statistics seen in numerical simulations could be explained as a superposition of these folded fields and Alfvén-like waves that propagate along the folds.

PACS numbers: 91.25.Cw, 47.27.Gs, 95.30.Qd, 47.27.Eq, 47.65.+a

arXiv:astro-ph/0308252 v1 14 Aug 2003

The term "MHD turbulence" embraces a number of turbulent regimes in conducting fluids. In this Letter, we consider what is perhaps the oldest MHD turbulence problem dating back to Batchelor's work in 1950 [1]: incompressible, randomly forced, nonhelical, homogeneous, isotropic MHD turbulence described by

$$\partial_t \mathbf{u} + \mathbf{u}\cdot\nabla\mathbf{u} = \nu\Delta\mathbf{u} - \nabla p + \mathbf{B}\cdot\nabla\mathbf{B} + \mathbf{f}, \qquad (1)$$

$$\partial_t \mathbf{B} + \mathbf{u}\cdot\nabla\mathbf{B} = \mathbf{B}\cdot\nabla\mathbf{u} + \eta\Delta\mathbf{B}, \qquad (2)$$

where the pressure p (determined from $\nabla\cdot\mathbf{u}=0$) and the magnetic field $\mathbf{B}$ are rescaled by ρ and $(4\pi\rho)^{1/2}$, respectively (ρ is density). Turbulence is excited by the random external forcing $\mathbf{f}$. No mean field is imposed. We are primarily interested in the case of large magnetic Prandtl number $\mathrm{Pr_m}=\nu/\eta$ which is appropriate for the warm ISM, coronal, and cluster plasmas [2]. Numerical evidence suggests that the popular choice $\mathrm{Pr_m}=1$ is in some respects similar to the large-$\mathrm{Pr_m}$ regime [3]. $\mathrm{Pr_m}\gg 1$ implies that the resistive scale $\ell_\eta \sim \mathrm{Pr_m}^{-1/2}\ell_\nu$ is much smaller than the viscous scale ℓ_ν. Thus, the problem has two scale ranges: the hydrodynamic (Kolmogorov) inertial range $\ell_0 \gg \ell \gg \ell_\nu \sim \mathrm{Re}^{-3/4}\ell_0$ (ℓ_0 is the forcing scale) and the subviscous range $\ell_\nu \gg \ell \gg \ell_\eta$.

For a moment, let us consider the traditional view of fully developed incompressible MHD turbulence in the presence of a strong, externally imposed mean field. This view is based on the idea of Iroshnikov [4] and Kraichnan [5] that it is a turbulence of strongly interacting Alfvén-wave packets. Their phenomenology, modified by Goldreich and Sridhar [6] to account for the anisotropy induced by the mean field, predicts steady-state spectra for magnetic and kinetic energies that are identical in the inertial range and have Kolmogorov $k^{-5/3}$ scaling. An essential feature of this description is that it implies scale-by-scale equipartition between magnetic and velocity fields: indeed, $\delta\mathbf{u_k} = \delta\mathbf{B_k}$ in an Alfvén wave. Numerics appear to confirm Alfvénic equipartition provided there is an imposed strong mean field $B_0 \gg u_{\mathrm{rms}}$ [7].

In the case of zero mean field, it has been widely assumed that essentially the same description applies, except now it is the large-scale magnetic fluctuations that play the role of effective mean field along which smaller-scale Alfvén waves can propagate. However, *numerical simulations of isotropic MHD turbulence do not show scale-by-scale equipartition between kinetic and magnetic energies.* There is a definite and very significant excess of magnetic energy at small scales. This is true for both $\mathrm{Pr_m}>1$ and $\mathrm{Pr_m}=1$ (Fig. 1). This result persists at the highest currently available resolution (1024^3, see [8]).

Let us consider the genesis of the magnetic field in isotropic MHD turbulence. As there is no mean field, all magnetic fields are fluctuations self-consistently generated by the small-scale dynamo. This type of dynamo is a fundamental mechanism that amplifies magnetic energy in sufficiently chaotic 3D flows with large enough magnetic Reynolds numbers and $\mathrm{Pr_m}\gtrsim 1$. The amplification is due to random stretching of the magnetic-field lines by the ambient velocity field. During the kinematic (weak-field) stage of the dynamo, the field energy grows exponentially in time and its spectrum is peaked at the resistive scale, $k_\eta \sim \mathrm{Pr_m}^{1/2}k_\nu$, and grows self-similarly [9, 10]. The dynamo growth rate is of the order of the turnover rate of the fastest eddies, which, in Kolmogorov turbulence, are the viscous-scale ones.

Although the bulk of the magnetic energy concentrates at the resistive scale, the dynamo-generated fields are not at all randomly tangled, but rather organized in folds within which the fields remain straight up to the scale of the flow and reverse direction at the resistive scale [11, 12]. One immediate implication of the folded



FIG. 1: Energy spectra in simulations with $\mathrm{Pr_m} = 1$, $\mathrm{Re}_\lambda \simeq 147$ and with $\mathrm{Pr_m} = 10$, $\mathrm{Re}_\lambda \simeq 43$. The thin (red) lines are the model-predicted spectra of the folded field component (normalized to have the same energy as the numerical spectra). The Kolmogorov slope is given for reference.

field structure is the criterion for the onset of nonlinearity: for incompressible MHD, the back reaction is controlled by the Lorentz tension force $\mathbf{B}\cdot\nabla\mathbf{B} \sim k_\parallel B^2$. This quantity depends on the parallel gradient of the field and does not know about direction reversals ($k_\parallel \sim k_\nu$ [12]). Balancing $\mathbf{B}\cdot\nabla\mathbf{B} \sim \mathbf{u}\cdot\nabla\mathbf{u}$, we find that back reaction is important when magnetic energy becomes comparable to the energy of the viscous-scale eddies. Clearly, some form of nonlinear suppression of the stretching motions at the viscous scale must then occur. However, the eddies at larger scales are still more energetic than the magnetic field and continue to stretch it at their (slower) turnover rate. When the field energy reaches the energy of these eddies, they are also suppressed and it is the turn of yet larger and slower eddies to exercise dominant stretching. The folded structure is preserved with folds elongating to the size ℓ_s of the dominant stretching eddy. The key question is whether ℓ_s can increase all the way to the outer scale or stabilizes just above the viscous scale [13].

The nonlinear suppression of stretching motions does not mean complete elimination of all turbulence: only the $\hat{\mathbf{b}}\hat{\mathbf{b}} : \nabla\mathbf{u}$ component of the velocity-gradient tensor leads to work being done against the Lorentz force and, therefore, must be suppressed. It is then natural to expect a local anisotropization of the velocity field. In this Letter, we demonstrate how a simple model accounting for this nonlinearly induced local anisotropy can produce solutions that are in remarkably good agreement with numerically observed magnetic-energy spectra.

The idea is to use the standard Kazantsev [9] model velocity, Gaussian and white in time, $\langle u^i(t,\mathbf{x})u^j(t',\mathbf{x}')\rangle = \delta(t-t')\kappa^{ij}(\mathbf{x}-\mathbf{x}')$, but let κ^{ij} depend on the local direction of the magnetic field, $\hat{\mathbf{b}} = \mathbf{B}/B$. In the Lagrangian frame (with local rotation transformed out), $\hat{\mathbf{b}}$ orients itself along the stretching Lyapunov direction of the flow, which stabilizes exponentially in time [14]. Therefore, in this frame, $\hat{\mathbf{b}}$ can be assumed to vary slowly with time. In the presence of one preferred direction defined by $\hat{b}^i\hat{b}^j$, the velocity correlator in $\mathbf{k}$ space has the following form

$$\begin{aligned}\kappa^{ij}(\mathbf{k}) \;=\; & \kappa^{(\mathrm{i})}(k,|\mu|)\big(\delta^{ij}-\hat{k}_i\hat{k}_j\big) + \kappa^{(\mathrm{a})}(k,|\mu|)\big(\hat{b}^i\hat{b}^j \\ & + \mu^2\hat{k}_i\hat{k}_j - \mu\hat{b}^i\hat{k}_j - \mu\hat{k}_i\hat{b}^j\big), \end{aligned} \tag{3}$$

where $\hat{\mathbf{k}} = \mathbf{k}/k$, $\mu = \hat{\mathbf{k}}\cdot\hat{\mathbf{b}}$. Let us ignore the spatial dependence of all quantities that vary at the flow scale and slower. The velocity will only enter via its gradient $u^i_j = \partial_j u^i$, which is now a function of time only with statistics $\langle u^i_m(t)u^j_n(t')\rangle = \delta(t-t')\int d^3k\, k_n k_m \kappa^{ij}(\mathbf{k})$. We can assume that $\hat{\mathbf{b}}$ also depends on time only, because it will always enter via the tensor $\hat{b}^i\hat{b}^j$, which varies at the scale of the flow (because of the folded structure of the magnetic field, the field's curvature is very small [12], so the fast spatial variation of $\hat{\mathbf{b}}$ is limited to sign reversals and cancels in $\hat{b}^i\hat{b}^j$). With these assumptions, the solution to Eq. (2) can be written in the form (cf. [15])

$$\mathbf{B}(t,\mathbf{x}) = \hat{\mathbf{b}}(t)\int d^3k_0\, \tilde{B}(t,\mathbf{k}_0)e^{i\mathbf{x}\cdot\mathbf{k}(t,\mathbf{k}_0)}, \tag{4}$$

where $\mathbf{k}(0,\mathbf{k}_0) = \mathbf{k}_0$ and

$$\partial_t\tilde{B} \;=\; \hat{b}^i\hat{b}^m u^i_m\tilde{B} - \eta k^2\tilde{B}, \tag{5}$$

$$\partial_t k_m \;=\; -u^i_m k_i, \tag{6}$$

$$\partial_t\hat{b}^i \;=\; \hat{b}^m u^i_m - \hat{b}^l\hat{b}^m u^l_m\hat{b}^i. \tag{7}$$

Equations (5–7) are a modification of the so-called zero-dimensional model of the dynamo [16]. A closed equations can be obtained for the joint PDF of $\tilde{B}$, $\mathbf{k}$, and $\hat{\mathbf{b}}$, $\mathcal{P}(\tilde{B},\mathbf{k},\hat{\mathbf{b}}) = \delta(|\hat{\mathbf{b}}|^2-1)\delta(\hat{\mathbf{b}}\cdot\mathbf{k})(4\pi^2k)^{-1}P(\tilde{B},k)$, via an averaging procedure analogous to, e.g., the one in Ref. [12]. The magnetic-energy spectrum $M(k) = (1/2)\int_0^\infty d\tilde{B}\,|\tilde{B}|^2P(\tilde{B},k)$ is then found to satisfy

$$\begin{aligned}\partial_t M \;=\; & \frac{1}{8}\gamma_\perp\frac{\partial}{\partial k}\bigg[(1+2\sigma_\parallel)k^2\frac{\partial M}{\partial k} \\ & - (1+4\sigma_\perp+10\sigma_\parallel)kM\bigg] \\ & + 2(\sigma_\perp+\sigma_\parallel)\gamma_\perp M - 2\eta k^2 M, \end{aligned} \tag{8}$$

where $\gamma_\perp = \int d^3k\, k_\perp^2\kappa_\perp$, $\sigma_\perp = (1/\gamma_\perp)\int d^3k\, k_\parallel^2\kappa_\perp$, $\sigma_\parallel = (1/\gamma_\perp)\int d^3k\, k_\parallel^2\kappa_\parallel$, $k_\perp = k(1-\mu^2)^{1/2}$, $k_\parallel = k\mu$, $\kappa_\perp = (1/2)\big(\delta^{ij}-\hat{b}^i\hat{b}^j\big)\kappa^{ij}$, $\kappa_\parallel = (1/2)\hat{b}^i\hat{b}^j\kappa^{ij}$, and κ^{ij} is defined in Eq. (3). In the isotropic case, $\kappa^{(\mathrm{i})} = \kappa^{(\mathrm{i})}(k)$, $\kappa^{(\mathrm{a})} = 0$, which gives $\sigma_\perp = 2/3$, $\sigma_\parallel = 1/6$. Equation (8) then

reduces to the standard equation for the magnetic-energy spectrum in the kinematic dynamo [9, 10]. With a zero-flux boundary condition imposed at low k [13], Eq. (8) has an eigenfunction (in the limit $\eta \to +0$)

$$M(k) \simeq k^s e^{\gamma t} K_0(k/k_\eta), \tag{9}$$

where K_0 is the Macdonald function, $k_\eta = \big[(1+2\sigma_\parallel)\gamma_\perp/16\eta\big]^{1/2}$, $s = 2(\sigma_\perp + 2\sigma_\parallel)/(1+2\sigma_\parallel)$, and $\gamma = (\gamma_\perp/8)\big[16(\sigma_\perp+\sigma_\parallel) - (1+2\sigma_\perp+6\sigma_\parallel)^2/(1+2\sigma_\parallel)\big]$. As magnetic back-reaction makes velocity more anisotropic, the values of $\sigma_\perp$, $\sigma_\parallel$ drop compared to the isotropic case, and so does the growth rate γ — until the dynamo is shut down (for a purely two-dimensional velocity, $\sigma_\perp = \sigma_\parallel = 0$ and $\gamma = -\gamma_\perp/8$). Thus, *saturation can be achieved purely by anisotropizing the statistics of the velocity field.*

How do we make connection from a theory based on the δ-correlated model velocity to the "real turbulence" that has a finite correlation time? The simplest prescription is to write finite expressions for equal-time velocity correlators by replacing the δ function by $1/\tau_{\rm c}$: $\langle u^i(\mathbf{k}) u^{j*}(\mathbf{k})\rangle = I^{ij}(\mathbf{k}) \sim \tau_{\rm c}^{-1}\kappa^{ij}(\mathbf{k})$. If the correlation time of a given type of motions is taken to be comparable to their turnover time, we can use $\tau_{\rm c} \sim \gamma_\perp^{-1}$ to write $\gamma_\perp = 3\,c_1\big[(2/5)\int d^3k\, k_\perp^2 I_\perp\big]^{1/2}$, where c_1 is a model-dependent constant of order unity. By the same token, $\sigma_\perp = \big[(2/3)\int d^3k\, k_\parallel^2 I_\perp / \int d^3k\, k_\perp^2 I_\perp\big]^{1/2}$ and $\sigma_\parallel = \big[(1/6)\int d^3k\, k_\parallel^2 I_\parallel / \int d^3k\, k_\perp^2 I_\perp\big]^{1/2}$.

In order to model gradual anisotropization of the velocity statistics by the nonlinear back reaction, we introduce the stretching wave number $k_{\rm s}(t)$ such that the total magnetic energy $W(t)$ at time t is approximately equal to the energy of the hydrodynamic eddies at $k > k_{\rm s}$ (before they feel the nonlinearity). We assume that the eddies at $k < k_{\rm s}$ remain isotropic (unaffected by back reaction), while those at $k > k_{\rm s}$ are two-dimensionalized. Specifically, for $k_0 < k < k_{\rm s}(t)$, let

$$4\pi k^2 I^{(\rm i)}(k,|\mu|) = E(k), \quad I^{(\rm a)}(k,|\mu|) = 0, \tag{10}$$

while for $k_{\rm s}(t) < k < k_\nu$,

$$4\pi k^2 I^{(\rm i)}(k,|\mu|) = 2r_{\rm 2D} E(k)\,\delta(\mu), \tag{11}$$
$$4\pi k^2 I^{(\rm a)}(k,|\mu|) = 2\tilde{E}(k)\,\delta(\mu). \tag{12}$$

Here $k_{\rm s}(t)$ is defined by $c_2\int_{k_{\rm s}(t)}^{k_\nu} E(k) = W(t)$ [$W(t)$ is the total magnetic energy at time t], $I^{(\rm i)}$ and $I^{(\rm a)}$ are coefficients of I^{ij} analogous to $\kappa^{(\rm i)}$ and $\kappa^{(\rm a)}$ [Eq. (3)], k_0 and k_ν are the forcing and viscous wave numbers, c_2 and $r_{\rm 2D}$ are adjustable parameters. We take $E(k) = C_{\rm K}\epsilon^{2/3}k^{-5/3}$ for $k_0 < k < k_\nu$. We emphasize that this is purely a choice of convenience: the behavior of our model is largely insensitive to the particular form of $E(k)$. $\tilde{E}(k)$ will not figure in what follows, because it multiplies μ in all relevant expressions. Coefficients in Eq. (8) now depend on $W(t)$: a straightforward calculation gives

$$\gamma_\perp(t) = \frac{6}{5}\bar{\gamma}\left[1 - \frac{1}{(1+W_0/W_\nu)^2}\right]^{-1/2}\big[\Gamma(t)\big]^{1/2}, \tag{13}$$

$$\sigma_\perp(t) = 4\sigma_\parallel(t) = \frac{2}{3}\left[\frac{1}{(1+W(t)/W_\nu)^2} - \frac{1}{(1+W_0/W_\nu)^2}\right]^{1/2}\big[\Gamma(t)\big]^{-1/2}, \tag{14}$$

$$\Gamma(t) = \frac{1}{(1+W(t)/W_\nu)^2} - \frac{1}{(1+W_0/W_\nu)^2} + \frac{5}{4}\,r_{\rm 2D}\left[1 - \frac{1}{(1+W(t)/W_\nu)^2}\right],$$

where $\bar{\gamma} = c_1\big[\int_{k_0}^{k_\nu} dk\, k^2 E(k)\big]^{1/2}$, the viscous-eddy energy is $W_\nu/c_2 = (3/2)C_{\rm K}\epsilon^{2/3}k_\nu^{-2/3}$, and the total energy of the velocity field (before suppression) is $W_0/c_2 = \int_{k_0}^{k_\nu} dk\, E(k)$. Equations (13–14) represent a generalization of the model first introduced in Ref. [13] and reduce to it when $r_{\rm 2D} = 0$. They include the effect of quasi-2D mixing of the folded magnetic fields by eddies whose stretching component has been suppressed. The spectrum of these mixing motions is modelled by Eq. (11), where $r_{\rm 2D}$ parametrizes the strength of the mixing relative to the original unsuppressed 3D turbulence.

The behavior of our model is easy to predict. The initial kinematic growth stage [$\gamma_\perp = (6/5)\bar{\gamma}$, $\sigma_\perp = 2/3$, $\sigma_\parallel = 1/6$, and $s = 3/2$, $\gamma = (3/4)\bar{\gamma}$ in Eq. (9)] lasts until the total magnetic energy becomes comparable to the energy of the viscous-scale eddies, $W \sim W_\nu$. After that, the velocity is gradually anisotropized, stretching is weakened, but mixing continues at $k > k_{\rm s}(t)$. A steady solution is reached as soon as $\sigma_\perp$ and $\sigma_\parallel$ have decreased enough to render $\gamma = 0$ in Eq. (9). This gives $\sigma_\perp = 4\sigma_\parallel \simeq 0.078$. The corresponding spectral exponent in the interval $k_\nu \ll k \ll k_\eta$ is $s \simeq 0.23$. This solution is valid in the limit $k_\eta \gg k_\nu$ ($\mathrm{Pr_m} \gg 1$), but convergence in $\mathrm{Pr_m}$ is only logarithmic. In practice, numerical solution of Eq. (8) with coefficients defined by Eqs. (13–14) shows that scale separations of about 7 decades are required for the scaling to be discernible. Obviously, this is not achievable in direct numerical simulations. We have, therefore, solved Eq. (8) numerically with physical parameters approximating those used in our direct numerical simulations of the MHD turbulence [3]. There are three adjustable constants: c_1, c_2, and $r_{\rm 2D}$. The results do not, in fact, depend very strongly on them: c_2 is irrelevant as it amounts to overall rescaling of energy, c_1 is a rescaling of time (though it also affects the value of k_η), and $r_{\rm 2D}$ has to vary by an order of magnitude to produce significant change. We have compared the model solutions for the *same* fixed values $c_1 = 1/3$ and $r_{\rm 2D} = 0.75$ with the (normalized) spectra obtained in numerical simulations at various $\mathrm{Pr_m}$ and Re. A sequence of runs with large $\mathrm{Pr_m}$ and low Re (the so-called viscosity-



FIG. 2: The thick lines are the normalized saturated energy spectra for simulations in the viscosity-dominated regime [3]. The thin (red) lines are the spectra predicted by our model.

dominated limit: a necessary compromise at current resolutions [3]) is very well fitted by the model in both kinematic (not shown) and nonlinear (Fig. 2) regimes (except at $k/2\pi = 1$, where boundary conditions are important).

It is extraordinary that a semi-heuristic minimal model has been able to reproduce non-asymptotic numerical spectral profiles so well. We do not claim that our model constitutes a quantitative theory of nonlinear dynamo. It does, however, provide a simple demonstration that the available numerical data (Fig. 2) is consistent with magnetic spectra that would exhibit a very flat positive spectral exponent in the interval $k_\nu \ll k \ll k_\eta$ if sufficiently large scale separations were resolved.

It is clear that the viscosity-dominated simulations (low Re) are described very well by our model. It is much harder to assess its performance for the case $\mathrm{Re} \gg 1$, $\mathrm{Pr_m} \gg 1$. If mixing by velocities in the interval $k_\mathrm{s} < k < k_\nu$ remains efficient [as is implied by our crude 2D approximation (11–12)], then k_s stabilizes at a value $\sim k_\nu$ and saturated magnetic energy scales with Re as the energy of the viscous eddies, $\langle B^2\rangle \sim \mathrm{Re}^{-1/2}\langle u^2\rangle$. This outcome does not appear to be borne out by the available numerical evidence, which rather suggests $\langle B^2\rangle \lesssim \langle u^2\rangle$ [8] (though limited resolutions do not allow one to make a definitive statement). In our own runs with $\mathrm{Pr_m} = 10$ and Taylor-microscale Reynolds number $\mathrm{Re}_\lambda \simeq 43$ and with $\mathrm{Pr_m} = 1$, $\mathrm{Re}_\lambda \simeq 147$, our model in its present form overestimates somewhat the magnetic energy at large k, but underestimates it at low k (Fig. 1): an indication of too much mixing in the model. Indeed, when $\mathrm{Re} \gg 1$, the nature of the anisotropized velocities in the interval $k_\mathrm{s} < k < k_\nu$ can be very different from the interchange-like motions that give the 2D mixing in the viscosity-dominated case. In Ref. [13], we argued that these motions are Alfvén waves that propagate along the folds. The saturated spectrum is then the result of a superposition of waves and folds (which accounts for the large amount of small-scale magnetic energy). Since the Alfvén waves are dissipated by viscosity, they can only exist if the stretching scale becomes much larger than the viscous scale: possibly as large as the outer scale. This is only allowed if the waves do not mix magnetic field as efficiently as the interchange motions do. For our model, the required modification would be that the mixing rate $\gamma_\perp$ should decrease with k_s. The dynamo saturation would then be due to a balance between stretching and mixing by partially anisotropized motions *at the stretching scale*.

Detecting the Alfvén waves along folds is a challenge for future numerical work. The main conclusion of the present study is that the nonlinear dynamo in a random one-scale flow can be described by a simple model where saturation is achieved via partial anisotropization of the ambient velocity, a result quantitatively supported by agreement with direct numerical simulations.

This work was supported by grants from PPARC (No. PPA/G/S/2002/00075), EPSRC (No. GR/R55344/01), UKAEA (No. QS06992). Simulations were done at UKAFF (Leicester) and NCSA (Illinois).

* Electronic address: a.schekochihin@imperial.ac.uk

[1] G. K. Batchelor, Proc. R. Soc. London A **201**, 405 (1950).
[2] L. M. Widrow, Rev. Mod. Phys. **74**, 775 (2002).
[3] A. A. Schekochihin, S. C. Cowley, S. F. Taylor, J. L. Maron, and J. C. McWilliams, Astrophys. J. (2003), submitted.
[4] P. S. Iroshnikov, Sov. Astron. **7**, 566 (1964).
[5] R. H. Kraichnan, Phys. Fluids **8**, 1385 (1965).
[6] P. Goldreich and S. Sridhar, Astrophys. J. **438**, 763 (1995).
[7] J. Maron and P. Goldreich, Astrophys. J. **554**, 1175 (2001).
[8] N. E. L. Haugen, A. Brandenburg, and W. Dobler, Phys. Rev. Lett. (2003), submitted [astro-ph/0303372].
[9] A. P. Kazantsev, Zh. Eksp. Teor. Fiz. **53**, 1806 (1967), [Sov. Phys.–JETP **26**, 1031 (1968)].
[10] R. M. Kulsrud and S. W. Anderson, Astrophys. J. **396**, 606 (1992).
[11] E. Ott, Phys. Plasmas **5**, 1636 (1998).
[12] A. Schekochihin, S. Cowley, J. Maron, and L. Malyshkin, Phys. Rev. E **65**, 016305 (2002).
[13] A. A. Schekochihin, S. C. Cowley, G. W. Hammett, J. L. Maron, and J. C. McWilliams, New J. Phys. **4**, 84 (2002).
[14] I. Goldhirsch, P.-L. Sulem, and S. A. Orszag, Physica D **27**, 311 (1987).
[15] Y. B. Zeldovich, A. A. Ruzmaikin, S. A. Molchanov, and D. D. Sokoloff, J. Fluid Mech. **144**, 1 (1984).
[16] A. Gruzinov, S. Cowley, and R. Sudan, Phys. Rev. Lett. **77**, 4342 (1996).