

Transport Predictions for Density-Gradient-Dominated Regimes

Erik M. Granstedt, G. W. Hammett

Princeton Plasma Physics Laboratory, Princeton, NJ

2011 U.S. Transport Task Force Workshop

Abstract

Tokamak energy confinement is often limited by turbulent transport due to temperature-gradient-driven instabilities; however, a low-recycling plasma boundary may support high edge temperatures, resulting in a flattened temperature profile and eliminating the drive mechanism for ITG and ETG turbulence. In this regime, modes driven by the density gradient may dominate the overall transport. To characterize transport in this regime, the dependence of particle and thermal fluxes as a function of density gradient with zero temperature gradient were examined, using the GYRO[Candy and Waltz, 2003] code. The nonlinear scaling of flux with density gradient indicates a nonlinear upshift in the critical gradient as reported elsewhere[Ernst et al., 2004]. In addition, the outward heat flux is less than the convective heat flux, due to preferential transport of low-energy particles. Thus, the transport attempts to drive a temperature gradient.

Abstract, contd.


The GYRO code has been used to assess the performance predictions for the Lithium Tokamak Experiment (LTX), a spherical torus designed to investigate the low-recycling lithium wall regime.[Majeski et al., 2009]

Linear simulations indicate that during active gas-puffing, both ITG and TEM modes are unstable. Charge-exchange losses with gas puff neutrals and weak electron-ion coupling suppress the edge T_i resulting in a steep ion temperature gradient and low T_i/T_e which drive the ITG mode.

Abstract, contd.

The relatively small size of LTX ($a/\rho_s \sim 25\text{--}40$) permits global nonlinear simulations of most of the plasma volume at reasonable computational cost, though it also complicates the simulations by making the role of the wall, sources, and sinks more important. Initial nonlinear simulations using predicted profiles during gas puffing failed to saturate indicating that the transport may be greater than expected. Saturation is achieved after relaxing the density and temperature profiles.

LTX Description



Cutaway view of internal shells and coils

Available Diagnostics:

- Comprehensive magnetics
- 2 mm scanning interferometer
- 1 mm fixed interferometer
- Multi-point Thomson Scattering
- Poloidal Lyman- α & Bolometer Arrays
- Edge Langmuir Probe
- Gas Puffer, Supersonic Gas Injector
- Molecular Cluster Injector
- Filterscopes & Visible survey spectrometers

Coming Soon:

- XUV spectrometer
- passive spectroscopy for n_i , T_i , u_i
- Digital Holography
- Neutral beam (late 2011 or early 2012)

LTX Achieved and Target Parameters

Parameter	Achieved (Ohmic)	Goal (beam-heated)
Major Radius (R_0)	40 cm	40 cm
Minor Radius (a)	26 cm	26 cm
Elongation (κ)	1.55	1.55
Toroidal Field (B_T)	1.8 kG	3.5 kG
Plasma Current (Ohmic) (I_p)	67 kA	> 250 kA
Central Density ($n_e(0)$)	$\sim 8 \times 10^{18} \text{ m}^{-3}$	$5 \times 10^{19} \text{ m}^{-3}$
Central T_e	$\sim 100 \text{ eV}$	> 500 eV
Central T_i	?	> 150 eV
Current Flattop	$\sim 5 \text{ ms}$ (20 ms duration)	> 100 ms

Reference Transport Model


- $D_{e,i} = \chi_i = \chi_e = \chi_i^{neo}$
- implemented in ASTRA with the equilibrium and stability code ESC
- flux boundary condition for the temperature profile
- ideal situation is zero recycling: no cold neutrals at the wall to suppress the edge temperature
- zero recycling is not compatible with edge particle fueling

Outline

- 1 Transport Model Linear Evaluation
 - During Gas Puffing
- 2 Parametric Scan: limit of $\nabla T_{e,i} = 0$
 - Linear Results
 - Nonlinear Results
- 3 LTX Global NL Simulations
 - Challenges
 - Preliminary Results
- 4 Conclusions

RTM Predictions during Gas Puffing

Sample low- β equilibrium ($\beta \sim 4\%$) [Majeski et al., 2009]




Objectives of gyro-kinetic simulations:

- develop interface between ASTRA/ESC and GYRO
- examine consistency of RTM with gyro-kinetic simulations
- gain insight into the expected density fluctuation levels in the experiment

$r/a = 0.7$, GYRO Linear γ during gas puffing


R_0/a	1.66
Δ	-0.11
κ	1.32
s_κ	0.018
δ	0.16
s_δ	0.20
q	1.12
s	1.38
ρ_*	0.035
ν_{ei}	0.022
ν_{ij}	0.008
a/L_n	1.56
a/L_{Te}	1.78
a/L_{Ti}	2.57
T_i/T_e	0.126
β_e	1.9%



Weak electromagnetic effects due to low β (2.1%). For $k_y \rho_s < 1.0$, TEM dominates; for $k_y \rho_s > 1.0$ ITG dominates.

$r/a = 0.7$, GYRO Linear ω_r during gas puffing


R_0/a	1.66
Δ	-0.11
κ	1.32
s_κ	0.018
δ	0.16
s_δ	0.20
q	1.12
s	1.38
ρ^*	0.035
ν_{ei}	0.022
ν_{ij}	0.008
a/L_n	1.56
a/L_{T_e}	1.78
a/L_{T_i}	2.57
T_i/T_e	0.126
β_e	1.9%



For $k_y \rho_s < 1.0$, drift is in the elec. diamag. direction; for $k_y \rho_s > 1.0$ in the ion direction.

$r/a = 0.8$, GYRO Linear γ during gas puffing


R_0/a	1.65
Δ	-0.17
κ	1.33
s_κ	0.11
δ	0.20
s_δ	0.38
q	1.44
s	2.48
ρ^*	0.029
ν_{ei}	0.026
ν_{ij}	0.011
a/L_n	2.80
a/L_{T_e}	1.98
a/L_{T_j}	3.24
T_i/T_e	0.111
β_e	1.0%



Strongly-driven TEM mode due to large density gradient.

$r/a = 0.8$, GYRO Linear ω_r during gas puffing

R_0/a	1.65
Δ	-0.17
κ	1.33
s_κ	0.11
δ	0.20
s_δ	0.38
q	1.44
s	2.48
ρ^*	0.029
ν_{ei}	0.026
ν_{ij}	0.011
a/L_n	2.80
a/L_{T_e}	1.98
a/L_{T_i}	3.24
T_i/T_e	0.111
β_e	1.0%



Shift to ion diamag. direction (ITG) for $k_y \rho_s > 2.2$.


Outline

- 1 Transport Model Linear Evaluation
 - During Gas Puffing
- 2 Parametric Scan: limit of $\nabla T_{e,i} = 0$
 - Linear Results
 - Nonlinear Results
- 3 LTX Global NL Simulations
 - Challenges
 - Preliminary Results
- 4 Conclusions

Linear, Local Parametric Scan

Values at $r/a = 0.75$ were taken, then $\nabla T \rightarrow 0$, $\rho_* \rightarrow 0.1\rho_*$ to get local limit of idealized flat $T_{e,i}$ regime. $T_i/T_e \rightarrow 1$ for reactor, low β (1.6%).


R_0/a	1.66
Δ	-0.14
κ	1.33
s_κ	0.06
δ	0.17
s_δ	0.28
q	1.25
s	1.95
ρ_*	0.0033
ν_{ei}	0.0
ν_{ij}	0.0



Linear, Local Parametric Scan


Real frequency can be in either electron or ion diamagnetic direction.

R_0/a	1.66
Δ	-0.14
κ	1.33
s_κ	0.06
δ	0.17
s_δ	0.28
q	1.25
s	1.95
ρ_*	0.0033
ν_{ei}	0.0
ν_{ij}	0.0



Linear, Local Parametric Scan


R_0/a	1.66
Δ	-0.14
κ	1.33
s_κ	0.06
δ	0.17
s_δ	0.28
q	1.25
s	1.95
ρ_*	0.0033
ν_{ei}	0.0
ν_{ij}	0.0



Small linear critical gradient: $a/L_n \approx 0.45$ ($R/L_n \approx 0.75$).

Linear, Local Parametric Scan


R_0/a	1.66
Δ	-0.14
κ	1.33
s_κ	0.06
δ	0.17
s_δ	0.28
q	1.25
s	1.95
ρ_*	0.0033
ν_{ei}	0.0
ν_{ij}	0.0



Outline

- 1 Transport Model Linear Evaluation
 - During Gas Puffing
- 2 Parametric Scan: limit of $\nabla T_{e,i} = 0$
 - Linear Results
 - Nonlinear Results
- 3 LTX Global NL Simulations
 - Challenges
 - Preliminary Results
- 4 Conclusions


Nonlinear, Local Parametric Scan



directory: "/p/gyro/egranste/sim/tem_ltx/nl/nuei0"Sun Apr 11 22:00:36 2010


Strong zonal flows
are driven
nonlinearly which
then damp the
turbulence resulting
in bursts of flux.

Nonlinear, Local Parametric Scan




Significant nonlinear
upshift due to zonal
flows.

Nonlinear, Local Parametric Scan





Flux is dominated
by modes with
 $k_y \rho_s < 0.5$.

Nonlinear, Local Parametric Scan





$Q_e, Q_i < 3/2\Gamma$: the transport is driving particles and heat outward but also redistributing free energy among ∇n , ∇T_e , and ∇T_i by driving a finite ∇T_e and ∇T_i .

Velocity-space fluxes



At low and moderate k_y/ρ_s which dominate the total flux ($k_y \rho_s \lesssim 0.6$), the flux is due to low-energy trapped particles.


Velocity-space fluxes, contd.



At higher k_y/ρ_s , passing and higher-energy particles diffuse radially outward, but these modes contribute very little to the total flux.



Variation with η_e

To test examine changes in the turbulence as ∇P is distributed between the ∇n and ∇T_e , nonlinear GYRO simulations were performed using “Cyclone Base Case” parameters with $\nabla T_i = 0$, $R_0/L_n + R_0/L_{T_e} = 20$ to test variation with η_e .




As η_e increases for fixed ∇P , and $\nabla T_i = 0$, $\delta T_e/T_e$ and Q_e increase, while $\delta n_e/n_e$, Γ , and Q_i decrease. Q_i remains $> 3/2\Gamma$. Relative RMS fluctuation levels behave similarly to the fluxes.

Variation with η_e , contd.



The relative contribution of various binormal wavenumbers remains relatively unchanged.

Variation with η_e , contd.



Cross-phase shows qualitative difference: for the dominant modes, δn_e and δT_e are nearly in-phase in the case of $\eta_e = 1$, while they are 90° – 180° out-of-phase for $\eta_e = 0$.

Outline

- 1 Transport Model Linear Evaluation
 - During Gas Puffing
- 2 Parametric Scan: limit of $\nabla T_{e,i} = 0$
 - Linear Results
 - Nonlinear Results
- 3 LTX Global NL Simulations
 - Challenges
 - Preliminary Results
- 4 Conclusions

LTX Global Nonlinear Simulation Challenges

LTX is relatively “small”: $a/\rho_s \sim 30$, **therefore:**

- must simulate most of radial cross-section
- LTX has no “local” limit; must do global simulation
- damping regions at boundaries of simulation domain reduce usable simulation volume

LTX Global Nonlinear Simulation Challenges

Convergence issues: (simulations centered at $r/a = 0.53$)

- $n = 1$ mode tends to blow-up
- for convergence: need to soften density gradient, adjust q profile
- simulations blow-up unless electrostatic, or T_e, T_i reduced by half
- converged simulations are unphysical

Convergence issues: (simulations centered at $r/a = 0.7$)

- T_e, T_i flattened to examine limiting case and eliminate ITG
- need to flatten density gradients at edge
- simulations blow-up unless strength of electromagnetic effects reduced by half


Outline

- 1 Transport Model Linear Evaluation
 - During Gas Puffing
- 2 Parametric Scan: limit of $\nabla T_{e,i} = 0$
 - Linear Results
 - Nonlinear Results
- 3 LTX Global NL Simulations
 - Challenges
 - Preliminary Results
- 4 Conclusions

Preliminary Results


- Limiting case of $\nabla T_{i,e} = 0$, centered at $r/a = 0.7$, β reduced by half. $n(r)$ flattened in the outer damping region. Large fluxes and significant density fluctuations across the radial domain.
- diffusion typically 10-30 times the RTM prediction
- large-scale mode extends across radial domain
- would result in flattening of density profiles
- predicted fluctuation level $\delta n / \bar{n}_e \sim 5-10\%$,
 $\delta n_e \sim 0.5-1 \times 10^{12} \text{cm}^{-3}$

Saturation of Fluxes




Large oscillations in the particle and heat fluxes. Small q_i/q_e primarily due to small T_i/T_e .

RMS Electron Density Fluctuations



Zero radial boundary condition may be affecting result. May need additional damping at boundaries.


Poloidal cut of ϕ fluctuations



Long,
radially-extended
eddies.

Midplane ϕ correlation function

potential $\langle C(\Delta r, \Delta \alpha, r_0 = 0.696) \rangle_{\alpha, t}$ with $182.900 < t < 36$



Likely not converged
with box.

Conclusions

- Linear simulations indicate that during active gas-puffing, both ITG and TEM modes are unstable. Due to a low T_i/T_e , the ITG peak is counter-intuitively at higher binormal wavenumber.
- Obtaining converged, global nonlinear gyro-kinetic simulations of a small device $a/\rho_s \sim 30$ is a challenge because of the boundary treatment effects and possibly a subcritical β instability.
- The RTM may be significantly under-predicting the particle and thermal transport.

Conclusions, contd.

- Nonlinear simulations with zero temperature gradients reveal a nonlinear upshift of the critical density gradient.
- The outward particle flux is primarily due to low-energy trapped particles interacting with modes with $k_y \rho_s \sim 0-0.6$ and being preferentially lost. Since $q_i, q_e < 3\Gamma/2$, the transport attempts to drive a finite temperature gradient.

Future Work

- Integrate a more complete transport model such as GLF23 or TGLF into ASTRA to better predict LTX performance.

Acknowledgments

The authors appreciate help from L. Zakharov with setting up ASTRA, and J. Candy and R. Waltz for use of the GYRO code. Work was performed under an NSF Graduate Research Fellowship. LTX is supported by US DOE contract DE-AC02-09CH11466.

References I

- J. Candy and R. Waltz, Journal of Computational Physics **186**, 545 (2003), ISSN 0021-9991.
- D. R. Ernst, P. T. Bonoli, P. J. Catto, W. Dorland, C. L. Fiore, R. S. Granetz, M. Greenwald, A. E. Hubbard, M. Porkolab, M. H. Redi, et al., Physics of Plasmas **11**, 2637 (2004).
- R. Majeski, L. Berzak, T. Gray, R. Kaita, T. Kozub, F. Levinton, D. Lundberg, J. Manickam, G. Pereverzev, K. Snieckus, et al., Nuclear Fusion **49**, 055014 (2009), ISSN 0029-5515.

Slides available at <http://princeton.edu/~erikg/>