

# Accurate calculation of Singular Current Densities in non-linear, ideal MHD equilibrium solutions

Max-Planck-Institut für Plasmaphysik (Dr. J. Loizu) & Princeton Plasma Physics Laboratory (Dr. S.R. Hudson)

[PROBLEM] For decades, singular current densities have been predicted in ideal 3D MHD equilibria,

- From  $\nabla \cdot \mathbf{j} = \nabla \cdot (\sigma \mathbf{B} + \mathbf{j}_\perp) = 0$ , where  $\mathbf{j}_\perp \equiv \mathbf{B} \times \nabla p / B^2$ , derive  $\mathbf{B} \cdot \nabla \sigma = -\nabla \cdot \mathbf{j}_\perp$ .
- Solution for  $\sigma = \frac{\mathbf{j} \cdot \mathbf{B}}{B^2}$ , assuming  $\mathbf{B} = \nabla \psi \times \nabla \theta + \iota(\psi) \nabla \zeta \times \nabla \psi$ , is  $\sigma_{m,n} = \underbrace{\frac{g_{m,n}(\psi) p'}{\iota m - n}}_{\text{Pfirsch-Schlüter}} + \underbrace{\Delta_{m,n} \delta(\psi - \psi_s)}_{\delta\text{-function}}$
- Conventional 3D, nonlinear, MHD equilibrium codes (such as NSTAB, VMEC) cannot resolve these currents because of numerical methods that assume continuous functions. . .

[IMPORTANCE] It is essential to resolve these singularities for (i) accurate equilibrium calculations, and these currents play a vital role in both (ii) linear stability theory *and* (iii) tearing mode theory.

[ADVANCEMENT] The STEPPED PRESSURE EQUILIBRIUM CODE (SPEC) [Hudson et al., PoP 19, 112502 (2012)] does allow for discontinuities and can resolve the singular currents!

Shown (below) is a sequence of equilibria [Loizu et al., PoP, 2015], where an island is successively “shielded” by ideal currents on so-called “ideal”-interfaces which are brought closer to the rational surface; and shown (right) is the resonant pressure-driven  $1/x$  current-density, [SPEC (red squares); analytic (black stars)], in excellent agreement.

