

The kink mode during the disruptions in tokamaks¹

Leonid E. Zakharov

Princeton Plasma Physics Laboratory, MS-27 P.O. Box 451, Princeton NJ 08543-0451

Theory Seminar, PPPL

January 10, 2008, Princeton, NJ

¹This work is supported by US DoE contract No. DE-AC020-76-CHO-3073.



Contents

1	Several slides from JET	4
2	The basics of the kink mode in the presence of the halo currents	10
2.1	Surface current due to plasma perturbation	11
2.2	Surface current and forces in the presence of the wet-zone	12
2.3	Vertical disruptions create the wet zone for the kink mode	14
2.4	Sideway forces and the thermal and current quenches	15
3	The theory of the Tokamak Kink Mode (TKM)	16
4	Potential implications of the kink mode theory	21
4.1	Ion saturation and Hiro currents	22
4.2	Ion collisionless skin depth and the Hiro currents	23
4.3	Where is the plasma edge located	26
5	Summary	29

The talk explains the locked $m/n = 1/1$ kink mode during the vertical disruption event when the plasma has an electrical contact with the plasma facing conducting surfaces. It is shown that the kink perturbation can be in equilibrium state even with a stable safety factor $q > 1$, if the halo currents, excited by the kink mode, can flow through the conducting structure. This suggests a new explanation of the toroidal asymmetry in magnetic measurements and so-called sideways forces on the in-vessel components during the disruption event.

In addition, the talk confirms the fundamental role of the halo currents (named here as "Hiro" currents), which interaction with the free boundary kink modes was in many occasions emphasized earlier by Hiro Takahashi and Eric Fredrickson. In fact, the physics of Hiro currents can explain four edge plasma stability regimes in tokamaks in a way consistent with DIII-D experiments.

1 Several slides from JET



Disruption Forces and Halo Currents in JET

December 2007

Peter de Vries, Mike Johnson, Fabien Lanoy, Valeria Riccardo



1 Disruption Forces and Halo Currents in JET – December 2007 - by Peter de Vries

EFDA JET **Two extreme cases**

- For ITER-like configuration and plasma current ($I_p=1.5\text{MA}$)
 - Same Δt_{QC} but large halo currents and large disruption force
 - Same Δt_{QC} but small halo currents and small disruption force

#57036 $\Delta t_{\text{QC}}=16\text{ms}$
 #57034 $\Delta t_{\text{QC}}=20\text{ms}$

#57036 $I_{\text{HALO}}=0.15\text{ [MA]}$
 #57034 $I_{\text{HALO}}=0.44\text{ [MA]}$

UKAEA Fusion

12 Disruption Forces and Halo Currents in JET – December 2007 - by Peter de Vries



EFDA JET **Halo Currents**

halo

plasma

vessel

UKAEA Fusion

4 Disruption Forces and Halo Currents in JET – December 2007 - by Peter de Vries





Introduction

- A disruption is a sudden (~20ms) loss of control and/or confinement of a tokamak plasmas.
 - Loss of vertical stability → Vertical Displacement Event (VDE)
 - Loss of confinement: energy quench → current quench (QC)

- The causes of disruptions are plentiful, often related to MHD stability or vertical stability.

- Disruptions could cause considerable damage to the structure of the tokamak device
 - Energy quench → large heat loads on in-vessel components
 - In JET up to 2MJ/m².
 - QC and VDE → Large electromagnetic forces on the surrounding structures.
 - In JET forces on the vacuum vessel up to 750Tonnes (7.5MN).

2 Disruption Forces and Halo Currents in JET – December 2007 - by Peter de Vries









Disruption forces on JET

- There is a large spread in measured disruption forces on JET
 - Even if normalised to I_p^2 .



Statistics of C1-C14 disruptions
 $N = N_0 \exp(-F/\delta F)$
 $N_0 = 400$
 $\delta F = 0.75$
 $N_{tot} = 1344$



Vessel force F_v [MN]
 $1.5MA$ $2.0MA$ I_p^2 [MA²] at $t_{DISR} - 0.2s$


5 Disruption Forces and Halo Currents in JET – December 2007 - by Peter de Vries





EFDA JET Halo currents and Forces

- Forces scale with the Halo currents on JET
 - Dominant source for the disruption forces on JET is due to halo currents
 - Although a distinct off-set is found (Coil forces? Eddy's?)
 - Halo currents (TPF x f) are smaller for higher q_{95} . [1]



[1] V. Riccardo, et al., PPCF 46 (2004) p925


7 Disruption Forces and Halo Currents in JET – December 2007 - by Peter de Vries



Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008

2 The basics of the kink mode in the presence of the halo currents

Deformation of the free plasma surface generates the surface current



$$\begin{aligned}
 r &= R - \rho \cos \omega, \\
 z &= \rho \sin \omega, \\
 \rho &= a + \xi(\omega, \varphi), \\
 q &= \frac{aB_\varphi}{RB_\omega},
 \end{aligned} \quad (2.1)$$

Its value is determined by the condition

$$\vec{B} \cdot \nabla \rho = 0. \quad (2.2)$$


If the plasma core is deformed in accordance with equilibrium conditions the only force acting on the plasma is the electromagnetic pressure

$$p_{j \times B} \nabla \rho = \vec{i} \times \vec{B}. \quad (2.3)$$

For stable plasma the pressure suppresses the perturbation




Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008



$$\mu_0 \vec{i} = -\xi_{11} \frac{2B_\phi}{R} \cos(\omega - \varphi) e_\varphi - \frac{1}{R} \xi_{11} \frac{2aB_\phi}{R^2} \sin(\omega - \varphi) e_\omega. \quad (2.4)$$

If $q > 1$ the kink mode 1/1 is stable (Shafranov, 1952)


2.2 Surface current and forces in the presence of the wet-zone



$$\vec{i}(\omega, \varphi) = -\frac{1}{a} I'_\omega \vec{e}_\varphi + \frac{1}{R} I'_\varphi \vec{e}_\omega, \quad (2.5)$$

$$F_x = \pi I_{pl} B_\phi (1 - q) \xi_{11},$$

Depending on the position of the wet-spot a troublesome sideways force can be generated



There is a finite size


$$\frac{\Delta\omega}{2\pi} = \frac{q-1}{q} \quad (2.6)$$

of the wet-zone for excitation of the kink mode.

If the wet-zone is self-generated, then there is a threshold for instability


2.3 Vertical disruptions create the wet zone for the kink mode

Kink-mode is linearly unstable in the case of axisymmetric wet-zone, e.g. at the vertical disruption events (VDE)



At the nonlinear stage the mode leads to a disbalance in plasma current measurements at different azimuths φ (TPF)

The force is applied to the wet-spot and is directed toward the plasma



Both thermal and current quench make the vertical equilibrium field

$$B_{z,ext} = -\frac{0.1I_{pl}}{R} \left(\ln \frac{8R}{a} + \beta_j + \frac{l_i}{2} - \frac{3}{2} \right), \quad \beta_j \rightarrow 0 \quad (2.7)$$

becomes excessive and pushes plasma inward, leading to the sideway force.

The trajectory of the plasma during VDEs depends on equilibrium



Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008

15

3 The theory of the Tokamak Kink Mode (TKM)

All the theory is reduced to relationship between ξ and i

For arbitrary m, n and perturbation of the form

$$\rho = a + \Re \left(\sum_{m,n} \xi_{mn} e^{im\omega - in\varphi} \right), \quad (3.1)$$

the surface currents are determined by the perturbed equilibrium theory (Zakharov, Sov.J. Plasma Physics,(1981), Reviews of Plasma Phys. v.11)

$$\begin{aligned} \vec{i} &= \nabla I(\omega, \varphi) \times \vec{e}_n = -\frac{1}{a} I'_\omega \vec{e}_\varphi + \frac{1}{R} I'_\varphi \vec{e}_\omega, \\ i(\omega, \varphi) &\equiv \Re \left(\sum_{m,n} i_{nm,\varphi} e^{im\omega - in\varphi} \right), \\ \mu_0 i_{nm,\varphi} &= \xi_{nm} \left[(m-1)j - \frac{2nB_\varphi}{R} \right] e^{im\omega - in\varphi} \end{aligned} \quad (3.2)$$

The electromagnetic pressure is given (for a uniform current density) by

$$p = 2I_{pl}B_\omega \Re \left(\sum_{nm} \xi_{nm} \frac{nq - m + 1}{m} (m - nq) e^{im\omega - in\varphi} \right), \quad (3.3)$$

Together with electro-dynamics of the vacuum vessel they are sufficient for simulation of disruptions




Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008

16

3 Theory of the Tokamak Kink Mode (TKM) (cont.)

Eq.(3.3) converts surface current into plasma displacement




$$\begin{aligned} \vec{i} &= \nabla I(\omega, \varphi) \times \vec{e}_n, \\ I(\omega, \varphi) &= I(x), \\ x &= x(\omega, \varphi), \\ \vec{i} &= -\frac{I'}{a} x'_\omega \vec{e}_\varphi + \frac{I'}{R} x'_\varphi \vec{e}_\omega \end{aligned} \quad (3.4)$$

The figure shows the current flow lines $x = \text{const}$ of the halo currents

3 Theory of the Tokamak Kink Mode (TKM) (cont.)

Eq.(3.3) converts surface current into plasma displacement




Even this hand-prescribed current profile reveals that

The shape of the surface displacement is consistent with the halo current supply in the wet-zone

3 Theory of the Tokamak Kink Mode (TKM) (cont.)

Theory of TKM should be linked with the electro-magnetic modeling



E.Lamzin Group, Applied. Math. Dept., STC "SINTEZ", Efremov Research Institute, St.Petersburg, RF




Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008

19

3 The theory of the Tokamak Kink Mode (TKM) (cont.)

The momentum $\simeq 2MN \cdot \text{sec}$ of the sideway force in ITER is equivalent to the hit of its VV by two 50 T T-90S tanks at the speed of 70 km/hour.



Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008

20

4 Potential implications of the kink mode theory

When applied to the quasi-stable plasmas, TKM stands for “Takahashi Kink Mode”

Hiro Takahashi and Eric Fredrickson revealed the apparent role of helical halo currents (“Hiro’s currents”) associated with the free boundary instabilities in DIII-D

4.1 Ion saturation and Hiro currents

Unlike in disruptions, in quasi-stationary plasmas the halo currents are limited by the ion-saturation current I_{ion}

The ion-saturation current is determined by the plasma particle flux to the wall

$$I_{ion,A} = 1.6 \cdot 10^{-19} \left. \frac{dN}{dt} \right|^{edge-wall},$$
$$\left. \frac{dN}{dt} \right|^{edge-wall} = \frac{1}{1 - R_{rec}} \cdot \left. \frac{dN}{dt} \right|^{core-edge} \simeq \frac{1}{1 - R_{rec}} \cdot \frac{N}{\tau_p}, \quad (4.1)$$
$$\frac{N}{\tau_p} \simeq 10^{22} \frac{1}{sec}$$

The Hiro currents I_H are limited by

$$I_H < I_{ion} \frac{\Delta_H}{\Delta_{SoL}}. \quad (4.2)$$

The Hiro currents introduce into stability theory two factors related to the plasma density

In collisionless SOL the Δ_H is related to the ion collisionless skin depth d_i

In collisionless SOL (with no emission from the plates) the ion inertia determines the Hiro current

$$\begin{aligned} \vec{B} &\equiv \nabla\psi \times \vec{e}_s, & \frac{e}{c} \cdot \frac{\partial\psi}{\partial t} &= m_i \frac{dV_i}{dt}, \\ enV_i &= \frac{1}{2}j_H = -\frac{1}{2} \cdot \frac{c}{4\pi} \Delta\psi, & & (4.3) \\ \frac{\partial}{\partial t} (\psi + d_i^{*2} \Delta\psi) &= 0_{resistive}, & d_i^* &\equiv \frac{d_i}{\sqrt{2}} = \frac{c}{\sqrt{2}\omega_{pi}} \end{aligned}$$

The $d_{i,cm}^*$ parameter determines the value of I_H and the plasma edge stability regime

The Hiro currents depth Δ_H is limited by

$$\Delta_H \simeq \min\{d_i, \Delta_{SOL}\} \quad (4.4)$$

4.2 Ion collisionless skin depth and the Hiro currents (cont.)

Four plasma edge stability regimes can be specified

1. The LiWF regime, $d_i^* \gg \Delta_{SOL}$. Negligible Hiro currents.

$$d_{i,cm}^* \simeq \frac{2.3}{\sqrt{n_{20}^{edge}}} \gg \Delta_{SOL} \quad (4.5)$$

Low recycling, low plasma density, no ELMs, no blobs, perfectly stable plasma edge. Exactly like in DIII-D experiments

2. The ELMy H-mode, $d_i^* \simeq \Delta_{SOL}$. Limited Hiro currents, reduced recycling. Resistive effects leads to relaxations.

3. L-mode, $d_i^* < \Delta_{SOL}$. Unlimited Hiro currents, total mess with blobs.

$$\pi i_H = \frac{I_{pl}}{a^2} \xi_{nm} (nq_{left} - nq) \quad (4.6)$$

4. High density disruptions, $d_i^* < \Delta_{SoL}$. Feed back between plasma core and the Hiro currents.

$$i_H \simeq \frac{I_{ion}}{2a} \simeq \frac{en^{edge} 4\pi^2 Ra}{2a} \gamma \xi, \quad (4.7)$$


$$2\pi^3 Ra \gamma e \cdot n^{edge} < \frac{I_{pl}}{a} (n_{qleft} - nq)$$

The plasma edge density explicitly enters into tokamak MHD stability condition.

4.3 Where is the plasma edge located

The mean free path, λ_D defines the position of the plasma edge


$$\lambda_{D,m} = 121 \frac{T_{keV}^2}{n_{20}} \quad (4.8)$$



At the plasma edge the temperatures are determined by particle fluxes

$$\frac{5}{2} \Gamma_{i,e}^{edge-wall} T_{i,e}^{edge} = \int_V P_{i,e} dV \quad (4.9)$$

The plasma edge is at the temperature pedestal



There is no confinement behind the plasma edge



Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008

27

ELMs, blobs, beginning of high density disruptions seem to be just the different types of Takahashi Kink Modes

At the same time, the TKMs, which are always unstable, are the primary candidate for determining the width of the edge temperature pedestal, where the confinement is destroyed.

Suppression of Hiro currents and TKM automatically leads to sharpening of the edge T-pedestal, in consistency with Todd's Evans observations in RPM experiments on DIII-D.

The long lasting misconception of "the edge transport barrier", based on naive static model of near-separatrix layer has nothing in common with the reality.

In fact, the so-called "edge transport barrier" is the
NO-confinement zone



Leonid E. Zakharov, Theory Seminar, PPPL, Princeton, NJ, January 10, 2008

28

1. The theory of the Tokamak Kink Mode explains the behavior of MHD instabilities in the REAL tokamak environment
2. Through the physics of Hiro currents, the edge plasma density was finally introduced into MHD theory as a decisive element of tokamak stability.

It is proved once again how consistent with the plasma physics and stability is the LiWF (delayed by 10 years in PPPL)

This regime can be described by the well-understood, ideal MHD model.



5 Summary (cont.)

At the same time, the understanding of stability in the conventional fusion can be illustrated by

