

*EXTENDED MHD MODELS*¹

J.D. Callen, C.C. Hegna, C.R. Sovinec
University of Wisconsin, Madison, WI 53706-1609

Fluid Closures Workshop, ORNL, Oak Ridge, TN, 22-24 March 2006

Theses:

- Most derivations of plasma MHD equations neglect dissipative effects (e.g., ideal MHD) or use collisional equations, closures (e.g., Braginskii).
- Extended MHD (ExMHD) equations developed from two-fluid equations with general closures for \vec{q} and $\overleftrightarrow{\pi}$ provide a reasonable basis for describing macroscopic plasmas — for arbitrary collisionality regimes along \vec{B} .
- Closures for \vec{q} and $\overleftrightarrow{\pi}$ are very anisotropic and must be developed with drives induced by $\vec{\nabla}n$, $\vec{\nabla}\vec{V}$, $\vec{\nabla}n$; different procedures should be used in developing and implementing parallel, cross and perpendicular closures.
- Micro-turbulence effects can be included, in principle, by averaging over higher mode numbers (\vec{k}) to add terms to the usual MHD-type equations.
- Extended MHD equations, closures for NIMROD have special requirements — write in terms of “primitive” vector fields \vec{V} , \vec{E} , \vec{B} ; in \vec{x} , t not \vec{k} , ω ; computable closures; plasma plus em fields conservation relations.

¹Research supported by DoE grant DE-FG02-86ER53218.

Outline

- **Extended MHD (ExMHD) model goals, assumptions**
- **Derivation Of ExMHD equations:**
 - Complete two-fluid equations
 - Extended MHD equations
 - Chapman-Enskog (C-E) procedure for neutral fluid closures
 - Anisotropic nature of closures
 - Approximations used in deriving collisional closures
- **Various types of closures for ExMHD:**
 - Moment approach — for Spitzer problem, Braginskii closures
 - Neoclassical-based closures
 - Complete set of ExMHD equations, specifications
 - Parallel kinetics — drift-kinetic C-E equation; PIC-based and “continuum” solutions
 - Perpendicular closures via fluid moments — diamagnetic/gyroviscosity, perpendicular
- **How can micro-turbulence effects be included?**
- **Special requirements for NIMROD simulations**
- **Summary**

Goal — of Extended MHD (ExMHD) Model

- The fundamental goals in developing an Extended MHD model are to:

Develop MHD-like equations that accurately model macroscopic plasma behavior in magnetized ($\omega_c \gg \nu$, $|\rho \vec{\nabla}| \ll 1$) plasmas — for analytics and initial value simulations,

Allow for arbitrary collisionality along the magnetic field \vec{B} — i.e., $|\lambda_e \nabla_{\parallel}| \gtrsim 1$ as well as the usual collisional (Braginskii) collisional regime ($|\lambda_e \nabla_{\parallel}| \ll 1$),

Incorporate any needed kinetic effects via closure relations that are obtained from moments of solutions of kinetic descriptions which are consistent with the extended MHD equations — i.e., that are obtained from a Chapman-Enskog-type procedure.

- Extended MHD equations should include the following MHD models:

Ideal MHD — MHD equations with no dissipation \implies isentropic equation of state,

Resistive MHD — MHD equations including dissipation due to plasma resistivity η ,

Reduced MHD – resistive MHD equations with compressional Alfvén waves removed,

Neoclassical MHD — MHD equations including poloidal flow damping, increased perpendicular inertia, and bootstrap current through parallel viscous forces on ions, electrons,

Electron and ion diamagnetic flow (“two-fluid”) effects — i.e., inclusion of ω_{*e} , ω_{*i} .

Assumptions For Extended MHD Equations, Simulations

- Some assumptions will be made to develop an Extended MHD model:

Plasma has evolved for many collision times before simulation begins so lowest order kinetic distribution is a Maxwellian. Equilibrium flows are assumed to be subsonic.

In $P_1(\vec{v}/v_T)$ moments, only flow and heat flow will be kept (neglect higher order flows).

Macroscopic instabilities evolve as plasma is driven slowly through instability threshold.

Classical, neoclassical and paleoclassical models provide minimum plasma transport.

Micro-turbulence is also in steady-state and representable by transport it induces via $\vec{\nabla} \cdot \vec{\Gamma}_a \simeq -\vec{\nabla} \cdot D_a \vec{\nabla}_\perp n$, etc. — assume \vec{k} -space of macro and micro instabilities separable.

Sources (e.g., heating) and sinks (e.g., neutrals) relevant only on transport time scale.

- These assumptions preclude considering the following physical processes:

Open field line regions — unless one adds appropriate parallel boundary conditions,

Velocity-space loss-cones near divertor separatrix — would need “direct-loss” terms,

Nonaxisymmetric effects of sources and sinks on the $t \lesssim 1/\nu \sim 1$ msec time scale,

Order unity pressure anisotropy — instead, $(p_\parallel - p_\perp)/p \sim \epsilon_\perp \ll 1$ is being assumed,

High \vec{k} , ω fluid modes causing 5D phase-space filamentation, nonlinear Landau damping,

FLR effects beyond second order (gyroviscosity with diamagnetic flows and \perp viscosity),

High mode numbers which require a kinetic description and yield micro-turbulence, e.g., $k_\theta \rho_S < 0.2$ limits toroidal mode numbers to $n \lesssim 0.2 r/q \rho_S \sim 20\text{--}50$ (ITER)?

Extended MHD (ExMHD) Model Has Some New Features

- The collisional friction force density \vec{R}_e is not just the resistivity but includes an electron heat flow \vec{q}_e — i.e., $\vec{R}_e = \frac{n_e e}{\sigma_0} \left(\vec{J} + \frac{3e}{5T_e} \vec{q}_e \right)$.
- The closure relations for \vec{q} and $\overleftrightarrow{\pi}$ (for both electrons and ions) are left in general form — to use different closures for different problems.
- The extreme anisotropy (parallel, cross, perpendicular to \vec{B}) in the closure relations is emphasized — to reflect different physics in each direction.
- Different procedures are proposed for obtaining parts of closure relations — kinetics for parallel but fluid for cross and perpendicular directions.
- An attempt is being made to include the dissipative, transport effects of micro-turbulence — via averaging over high mode number processes and assuming they can be separated from the macroscopic (ExMHD) processes.
- The ExMHD model facilitates a more precise assessment of the range of validity of MHD-type models — when there is no significant entropy production, or with selected entropy production mechanisms (e.g., resistivity).

Fluid Moment Equations From Plasma Kinetic Equation

- The rigorous Plasma Kinetic Equation (PKE) to begin from is

$$\boxed{\frac{\partial f}{\partial t} + \vec{v} \cdot \vec{\nabla} f + \frac{q}{m} (\vec{E} + \vec{v} \times \vec{B}) \cdot \vec{\nabla}_v f = \mathcal{C}\{f\}.}$$

- Exact fluid moment equations for each plasma species result from velocity-space moments ($\int d^3v \vec{v}^n$, $n = 0, 1, 2$) of this fundamental kinetic equation:

$$\begin{aligned} n = 0, |\vec{v}|^0, \text{ density} \quad & \frac{\partial n}{\partial t} + \vec{\nabla} \cdot n \vec{V} = 0, \quad \{\vec{\nabla} \vec{V}\} \equiv \frac{1}{2} [\vec{\nabla} \vec{V} + (\vec{\nabla} \vec{V})^T] - \frac{1}{3} \overleftrightarrow{\mathbf{I}} (\vec{\nabla} \cdot \vec{V}), \\ n = 1, \vec{v}, \text{ momentum} \quad & mn \frac{d\vec{V}}{dt} = nq(\vec{E} + \vec{V} \times \vec{B}) - \vec{\nabla} p - \vec{\nabla} \cdot \overleftrightarrow{\pi} + \vec{R}, \quad \frac{d}{dt} \equiv \frac{\partial}{\partial t} + \vec{V} \cdot \vec{\nabla}, \\ n = 2, v^2, \text{ energy} \quad & \frac{3}{2} n \frac{dT}{dt} + nT \vec{\nabla} \cdot \vec{V} = -\vec{\nabla} \cdot \vec{q} - \overleftrightarrow{\pi} : \{\vec{\nabla} \vec{V}\} + Q, \quad p \equiv nT, \\ \text{or entropy} \quad & \frac{\partial(ns)}{\partial t} + \vec{\nabla} \cdot \left(ns \vec{V} + \frac{\vec{q}}{T} \right) = \frac{1}{T} (-\vec{q} \cdot \vec{\nabla} \ln T - \overleftrightarrow{\pi} : \{\vec{\nabla} \vec{V}\} + Q), \quad s \equiv \ln(T^{3/2}/n). \end{aligned}$$

- These moment equations need closure moments for \vec{q} and $\overleftrightarrow{\pi}$ ($\vec{v}_r \equiv \vec{v} - \vec{V}$):

$$\text{heat flux } \vec{q} \equiv \int d^3v \vec{v}_r \left(\frac{mv_r^2}{2} - \frac{5}{2} \right) f, \quad \text{stress tensor } \overleftrightarrow{\pi} \equiv \int d^3v m \left(\vec{v}_r \vec{v}_r - \frac{v_r^2}{3} \overleftrightarrow{\mathbf{I}} \right) f.$$

Definitions Of Relevant Velocity-Space Integrals

- Fluid moments in terms of velocity-space integrals of distribution function:

density: $n(\vec{x}, t) \equiv \int d^3v f,$

flow velocity: $\vec{V}(\vec{x}, t) \equiv \frac{1}{n} \int d^3v \vec{v} f,$

temperature: $T(\vec{x}, t) \equiv \frac{1}{n} \int d^3v \frac{m|\vec{v}_r|^2}{3} f, \quad \vec{v}_r \equiv \vec{v} - \vec{V}(\vec{x}, t).$

- Velocity-space moments of the Coulomb collision operator:

density: $0 \equiv \int d^3v \mathcal{C}\{f\},$

momentum: $\vec{R}_e \equiv \int d^3v m_e \vec{v} \mathcal{C}\{f\} = -m_e n_e \nu_e [(\vec{V}_e - \vec{V}_i) - \alpha \vec{q}_e / n_e T_e + \dots] \sim n_e e \eta \vec{J},$

energy: $Q_e \equiv \int d^3v \frac{m_e |\vec{v}_r|^2}{2} \mathcal{C}\{f\} = -\frac{3}{2} n_e \nu_e \frac{m_e}{m_i} (T_e - T_i) \underbrace{-\vec{R}_e \cdot (\vec{V}_e - \vec{V}_i)}_{\sim +\eta J^2}.$

- Note that closure moment for \vec{q}_e is needed to specify frictional force \vec{R}_e and energy transfer Q_e . Need parallel electron heat flow for Spitzer resistivity. (Braginskii high collisionality closure writes \vec{q}_e in terms of E_{\parallel} and $\nabla_{\parallel} T_e$.)

Collisional Closures Deduced Via Chapman-Enskog Approach

- In a neutral fluid, assuming the lowest order kinetic equation is dominated by the collision operator (i.e., $\mathcal{C}\{f_0\} \simeq 0$), the lowest order solution $f_0 = f_M$ is a “dynamic” Maxwellian with “parameters” $n(\vec{x}, t)$, $T(\vec{x}, t)$, $\vec{V}(\vec{x}, t)$:

$$f_M(\vec{x}, \vec{v}, t) = n(\vec{x}, t) \left(\frac{m}{2\pi T(\vec{x}, t)} \right)^{3/2} e^{-m|\vec{v}_r|^2/2T(\vec{x}, t)}, \quad \vec{v}_r \equiv \vec{v} - \vec{V}(\vec{x}, t).$$

- Chapman-Enskog procedure: Next order equation is obtained by substituting $f = f_M + \delta f$ into kinetic equation, making use of density, momentum and energy conservation equations to remove dependences on $\partial n/\partial t$, $\partial \vec{V}/\partial t$ and $\partial T/\partial t$, and neglecting higher order corrections ($\sim 1/\nu$):

$$\mathcal{C}\{\delta f\} \simeq \left[\left(\frac{m|\vec{v}_r|^2}{2T} - \frac{5}{2} \right) \vec{v}_r \cdot \vec{\nabla} \ln T + \frac{m}{T} \{ \vec{\nabla} \vec{V} \} : \left((\vec{v}_r \vec{v}_r - \frac{|\vec{v}_r|^2}{3} \mathbf{I}) \right) \right] f_M \sim -\nu \delta f.$$

- Inverting collision operator yields δf whose velocity-space moments provide needed closure relations (collision length $\lambda \equiv v_T/\nu$, $v_T \equiv \sqrt{2T/m}$):

$$\vec{q} = -\kappa^m \vec{\nabla} T = -n \chi^m \vec{\nabla} T, \text{ with “molecular” heat diffusivity } \chi^m \sim v_T^2/\nu = \nu \lambda^2,$$

$$\vec{\pi} = -2\nu^m \{ \vec{\nabla} \vec{V} \} = -nm \mu^m \{ \vec{\nabla} \vec{V} \}, \text{ “molecular” viscosity } \mu^m \sim \nu \lambda^2, \vec{\nabla} \cdot \vec{\pi} \simeq -\nu^m \nabla^2 \vec{V}.$$

Magnetized Plasmas Are Very Anisotropic ($\parallel, \wedge \perp$, to \vec{B})

- Braginskii [1] used a Chapman-Enskog procedure and an ordering scheme for magnetized ($\omega_c \equiv qB/m \gg \nu$), collisional ($\nu \gg \omega, k_{\parallel} v_T$) plasmas:

$$\perp \text{ to } \vec{B}: \text{ small gyroradius, } \varrho \equiv v_T/\omega_c \implies \epsilon_{\perp} \sim |\varrho \vec{\nabla}_{\perp}| \ll 1.$$

$$\parallel \text{ to } \vec{B}: \text{ short collision length, } \lambda \equiv v_t/\nu \implies \epsilon_{\parallel} \sim |\lambda \nabla_{\parallel}| \ll 1.$$

- Conductive heat flux closure moment is found to have parallel (\parallel), cross (\wedge , in flux surface) and perpendicular (\perp , across flux surfaces) components:

$$\vec{q} = -n \chi_{\parallel} \nabla_{\parallel} T - n \chi_{\wedge} (\vec{B}/B) \times \vec{\nabla} T - n \chi_{\perp} \vec{\nabla}_{\perp} T, \quad \text{in which } \vec{\nabla}_{\perp} \equiv - (1/B^2) \vec{B} \times (\vec{B} \times \vec{\nabla}),$$

$$\text{parallel heat conduction: } \chi_{\parallel} \sim \nu \lambda^2 \sim \epsilon_{\parallel}^2 \epsilon_{\perp}^0 \implies \text{fast } (t \sim 1/\nu), \parallel T \text{ equilibration,}$$

$$\text{cross (diamagnetic heat flow): } \chi_{\wedge} \sim v_T \varrho \sim \epsilon_{\perp} \implies \text{slower, diamag. flows in surface,}$$

$$\text{perpendicular heat conduction: } \chi_{\perp} \sim \nu \varrho^2 \sim \epsilon_{\perp}^2 \implies \text{slowest, radial heat transport.}$$

- Stress tensor has similar form: $\overleftrightarrow{\pi} = \overleftrightarrow{\pi}_{\parallel} + \overleftrightarrow{\pi}_{\wedge} + \overleftrightarrow{\pi}_{\perp}$ with similar scalings $\overleftrightarrow{\pi}_{\parallel} \sim \epsilon_{\parallel}^2 \epsilon_{\perp}^0$ (parallel stress), $\overleftrightarrow{\pi}_{\wedge} \sim \epsilon_{\perp}$ (gyroviscosity) and $\overleftrightarrow{\pi}_{\perp} \sim \epsilon_{\perp}^2$ (\perp visc.).

[1] S.I. Braginskii, in *Reviews of Plasma Physics* (Consultants Bureau, NY, 1965), Vol I, p 205.

Comments On Collisional Magnetized Plasma Equations

- The collisional fluid equations use these anisotropic closures and the resultant “two-fluid” equations are known as the Braginskii equations.
- Braginskii closures and equations are derived using the following major approximations, which determine their range of validity:

short collision length, $\epsilon_{\parallel} \sim \lambda \nabla_{\parallel} \ll 1$ — not valid for most tokamak plasma regimes,
small gyroradius, $\epsilon_{\perp} \sim \rho \vec{\nabla}_{\perp} \ll 1$ — equil. ok, but need $k_{\perp} \rho \ll 1$ for perturbations,
slow processes, $\partial/\partial t \ll \nu$ — equilibrium ok, but need $\omega/\nu \ll 1$ for perturbations,
negligible anomalous transport — add transport coefficients from microturbulence?

- Critiques of the Braginskii equations:

They neglect effects due to collisions with neutrals or energetic (e.g., fast ion) particles — but these transport-time-scale (slow) effects can mostly just be added as “sources.”

They do not include direct loss processes (e.g., near separatrix, on open field lines).

The stress tensor $\overleftrightarrow{\pi}$ is driven not just by $\{\vec{\nabla}\vec{V}\}$ but also by a comparable $\{\vec{\nabla}\vec{q}\}$ [2].

[2] A.B. Mikhailovskii, *Theory of Plasma Instabilities* (Atomdat, Moscow, 1977), Vol 2, p 307-325 (in Russian);
A.B. Mikhailovskii and V.S. Tsypin, *Plasma Physics* 13, 785 (1971); *ibid.*, *Beitr. Plasmaphys.* 24, 335 (1984).

Extended MHD Model Derived From Two-Fluid Equations

- Assume for the moment that anisotropic closures for \vec{q} and $\overleftrightarrow{\pi}$ can be obtained for both electrons and ions for relevant situations.
- Then, adding, subtracting electron and ion density and momentum equations one obtains general “Extended MHD” equations:

$$\begin{array}{ll}
 \text{density} & \frac{\partial \rho_m}{\partial t} + \vec{\nabla} \cdot \rho_m \vec{V} = 0, \quad \rho_m \equiv \frac{\sum_s n_s m_s}{\sum_s m_s} \simeq n_i, \quad \vec{V} \equiv \frac{\sum_s n_s m_s \vec{V}_s}{\sum_s n_s m_s} \simeq \vec{V}_i, \\
 \text{charge density} & \vec{\nabla} \cdot \vec{J} = 0, \quad \vec{J} \equiv e(n_i Z_i \vec{V}_i - n_e \vec{V}_e), \\
 \text{momentum} & \rho_m \frac{d\vec{V}}{dt} = \vec{J} \times \vec{B} - \vec{\nabla} P - \vec{\nabla} \cdot \overleftrightarrow{\Pi}, \quad P \equiv p_e + p_i, \quad \overleftrightarrow{\Pi} \simeq \overleftrightarrow{\pi}_i + \overleftrightarrow{\pi}_e \simeq \overleftrightarrow{\pi}_i, \\
 \text{Ohm's law} & \vec{E} + \vec{V} \times \vec{B} = \underbrace{\frac{\vec{R}_e}{n_e e}}_{\sim \eta \vec{J}} + \underbrace{\frac{\vec{J} \times \vec{B} - \vec{\nabla} p_e - \vec{\nabla} \cdot \overleftrightarrow{\pi}_e}{n_e e}}_{\text{Hall terms}} + \underbrace{\frac{m_e}{e^2} \frac{d}{dt} \left(\frac{\vec{J}}{n_e} \right)}_{\text{electron inertia}}.
 \end{array}$$

- Main effects of closures come in parallel Ohm's law and equation of state for the total plasma pressure P obtained from plasma entropy evolution:

$$\frac{d}{dt} \left(\ln \frac{P}{\rho_m^\Gamma} \right) = \frac{\Gamma - 1}{P} \left(p_e \frac{ds_e}{dt} + p_i \frac{ds_i}{dt} \right) \simeq \frac{\Gamma - 1}{P} \left(- \underbrace{\vec{\nabla} \cdot \vec{q}_e}_{\sim \nu_e \epsilon_{\parallel}^2} - \underbrace{\{\vec{\nabla} \cdot \vec{V}_i\}}_{\sim \nu_i \epsilon_{\perp}^2} : \overleftrightarrow{\pi}_i + \underbrace{\eta J^2}_{\sim 1/\tau_E} \right), \quad \Gamma \equiv \frac{5}{3}.$$

Comments On Extended MHD Equations

- Extended MHD equations derived from general two-fluid equations provide a useful, exact formal framework for MHD-type simulations.
- Natural variables to be advanced are usual MHD ones: $\rho_m, \vec{V}, \vec{J} \rightarrow \vec{B}, P$.
- Collisional friction force \vec{R}_e and closure relations for \vec{q} and $\overleftrightarrow{\pi}$ should be left unspecified, but split into parallel, cross and perpendicular parts:
 - parallel from C-E based kinetic analysis — continuum or PIC type?, involves $V_{\parallel}, q_{\parallel}, T$,
 - cross (gyroviscosity) — from fluid-type response, to sufficient accuracy,
 - perpendicular — from collisional effects on gyroviscosity-influenced flows in surface.
- Because plasma is Maxwellian to lowest order, the collisional entropy defined by $s = \ln(T^{3/2}/n)$ is a relevant quantity for each plasma species.
- Plasma pressure evolution should be obtained from entropy evolution:

$$\frac{d}{dt} \left(\ln \frac{P}{\rho_m^{\Gamma}} \right) = \frac{\Gamma - 1}{P} \left(p_e \frac{ds_e}{dt} + p_i \frac{ds_i}{dt} \right) \simeq \frac{\Gamma - 1}{P} \left(- \underbrace{\vec{\nabla} \cdot \vec{q}_e}_{\sim \nu_e \epsilon_{\parallel}^2} - \underbrace{\{\vec{\nabla} \vec{V}_i\} : \overleftrightarrow{\pi}_i}_{\sim \nu_i \epsilon_{\perp}^2} + \underbrace{\eta J^2}_{\sim 1/\tau_E} \right), \quad \Gamma \equiv \frac{5}{3}$$

Comments On Closures For Extended MHD Equations

- The only general, analytic closures are the collisional (Braginskii) ones.
- The main limitation in using Braginskii closures is the high collisionality requirement for the parallel kinetics: $\epsilon_{\parallel} \sim (v_T/\nu)\nabla_{\parallel} = \lambda\nabla_{\parallel} \ll 1$.
- The closures should be determined from a Chapman-Enskog-type procedure so the kinetics used to obtain them does not produce “extra” $\partial\delta n/\partial t$, $\partial\delta\vec{V}/\partial t$, and/or $\partial\delta T/\partial t$ contributions to the equations:
The usual drift-kinetic and gyro-kinetic equations are not developed using a Chapman-Enskog-like procedure and hence usually produce δn , $\delta\vec{V}$, and/or δT terms. Formal Chapman-Enskog-type procedures and resultant drift-kinetic equations have been developed for arbitrary \parallel collisionality [3-5], but they are rather complicated.
- The anisotropic components of the closures can be handled differently:
parallel: in general a kinetic analysis must be used, including collisional effects,
cross: fluid-type analysis, gyroviscosity for these diamagnetic flow type effects,
perpendicular: fluidlike radial transport due to collisional effects on diamagnetic flows.

[3] K.C. Shaing and D.A. Spong, Phys. Fluids B 2, 1190 (1990) — first Chapman-Enskog-like formalism.

[4] J.P. Wang and J.D. Callen, Phys. Fluids B 4, 1139 (1992) — axisymmetric \vec{B} , neoclassical formalism.

[5] Z. Chang and J.D. Callen, Phys. Fluids B 4 1167 (1992) — sheared slab model, with Landau damping.

Comments On Closures For Extended MHD (continued)

- Friction forces \vec{R} and stress tensors $\overleftrightarrow{\pi}$ are most fundamentally, generally written in terms of \vec{V} and \vec{q} — rather than \vec{V} and $\vec{\nabla}T$ Braginskii uses.
- The parallel Ohm's law is governed experimentally by the neoclassical Ohm's law and apparently not affected [6,7] by microturbulence — because $k_{\parallel} \ll k_{\perp}$ and hence their parallel momentum transfer is small.
- Temporal regimes — it seems there are two MHD regimes of interest:
 - “fast MHD” ($\omega \gg \nu$) — little entropy production, closures not very important?
 - “slow MHD” ($\omega \ll \nu$) — collision-dominated closures and dissipation critical.
- Spatial regimes — very anisotropic and different physics each direction:
 - parallel: need more general kinetic-based formalism, closures for $k_{\parallel}\lambda \sim 1$
 - cross (in flux surface): need separation from drift-wave-type microturbulence
 - $\implies k_{\theta}\rho_S < 0.2? \implies$ toroidal mode numbers $n \lesssim 0.3 r/(q \rho_S) \sim 20\text{--}50(\text{ITER})?$
 - perpendicular (across flux surface): avoid FLR effects on resistive layer widths
 - $\implies k_x \rho_i < 1$ with $\delta_{\eta} \sim r/(mS)^{1/3} \implies nS \lesssim (r/\rho_i)^3/q \sim 10^6\text{--}3 \times 10^7(\text{ITER})?$

[6] K.C. Shaing, Phys. Fluids 31, 8 (1988) — for electrostatic microturbulence.

[7] F.L. Hinton, R.E. Waltz, and J. Candy, Phys. Plasmas 11, 2433 (2004) — including $\delta\vec{B}_{\perp}$ effects.

Moment Expansion Solution Of “Kinetic” Spitzer Problem

- Electron flow, current induced by electric field is called Spitzer problem:

$$\frac{q_e}{m_e} \vec{E} \cdot \frac{\partial f_M}{\partial \vec{v}} = \mathcal{C}\{\delta f\} \implies \delta f = -\mathcal{C}^{-1} \left\{ \frac{q_e \vec{v} \cdot \vec{E}}{T} f_M \right\} \implies \vec{J} = q_e \int d^3v \vec{v} \delta f \equiv \sigma_{\text{Sp}} \vec{E}.$$

- In moment approach one takes $\int d^3v \vec{v} L_i^{(3/2)}$ moments of kinetic equation and obtains a matrix equation to be solved for \vec{V} , \vec{q} , etc. induced by \vec{E} :

$$n_e e \begin{pmatrix} \vec{E} \\ 0 \\ \vdots \end{pmatrix} = - \frac{m_e n_e \nu_e}{Z} \underbrace{\begin{pmatrix} \ell_{00} & \ell_{01} & \cdots \\ \ell_{10} & \ell_{11} & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix}}_{L_{ij}} \begin{pmatrix} \vec{V}_e - \vec{V}_i \\ -\frac{2}{5n_e T_e} \vec{q}_e \\ \vdots \end{pmatrix} \implies \begin{cases} \vec{J} \equiv -n_e e (\vec{V}_e - \vec{V}_i) \\ = \frac{n_e e^2}{m_e \nu_e} Z [L_{ij}^{-1}]_{00} \vec{E}. \end{cases}$$

- One can show [8] that inverting the friction matrix L_{ij} yields variational solution of Spitzer problem and hence plasma electrical conductivity σ :

1 × 1 matrix inversion yields $\sigma_0 \equiv \frac{n_e e^2}{m_e \nu_e}$, which is reference (\perp) conductivity,

2 × 2 matrix inversion yields $\sigma_{\text{Sp}} = \frac{1}{\alpha_e} \sigma_0$, $\alpha_e = \frac{\sqrt{2+Z}}{\sqrt{2+13Z/4}} \lesssim 5\%$ accuracy ($< \frac{1}{\ln \Lambda} \sim 0.07$),

3 × 3 matrix inversion yields $\sigma_{\text{Sp}} = \frac{1}{\alpha_e} \sigma_0$, with $\lesssim 1\%$ accuracy in α_e ($\simeq 0.51$ for $Z=1$).

[8] S.P. Hirshman, 21, 1295 (1978) — variational solution of Spitzer problem via moments.

Comments On Moment Approach Solutions Of Kinetics

- Moment approach matrix solution of Spitzer problem also produces electron heat flux induced by the electric field, $\frac{2}{5n_e T_e} \vec{q}_e = \frac{Ze}{m_e \nu_e} [L_{ij}^{-1}]_{01} \vec{E}$.

This is a key contribution to Spitzer conductivity (with $\geq 2 \times 2$ matrix inversion) since it converts \parallel friction force $R_{e\parallel}$ from reference (\perp) to \parallel Spitzer electrical conductivity:

$$\vec{R}_e = -m_e n_e \nu_e \left[(\vec{V}_e - \vec{V}_i) - \frac{3}{5n_e T_e} \vec{q} \right] = \frac{n_e e}{\sigma_0} \left(\vec{J} - \frac{9Z/4}{\sqrt{2+Z}} \sigma_0 \vec{E} \right) \implies \vec{J} = \sigma_0 \left(\frac{\sqrt{2+13Z/4}}{\sqrt{2+Z}} \right) \vec{E} = \sigma_{\text{Sp}} \vec{E}.$$

- In general there are additional “thermodynamic” drives (beyond the electric field \vec{E}) due to $\vec{\nabla} \ln p$ and $\vec{\nabla} \ln T \implies$ transport fluxes $\vec{\Gamma}, \vec{q}$.

- Braginskii collisional closures were obtained using moment approach:

Effects of all “forces” ($\vec{E}, \vec{\nabla} \ln p, \vec{\nabla} \ln T, \{\vec{\nabla} \vec{V}\}$) were determined simultaneously \implies Onsager symmetry, thermal force effect ($0.71 \nabla_{\parallel} T_e$), Ettinghausen effect, etc.;

4×4 matrix inversion was used for accurate numerical coefficients;

However, really only need 2×2 approach for order $1/\ln \Lambda \sim 5\%$ accuracy in resistivity but factor of 2 accuracy in thermal diffusivity χ — need 3×3 for similarly accurate χ .

- Moment approach shows that, at least in the collisional regime, the relevant fluid moment variables are $(n, T), (\vec{V}, \vec{q}), (\overleftrightarrow{\pi}, \overleftrightarrow{\pi}_q)$, with closures for $\vec{q}, \overleftrightarrow{\pi}, \overleftrightarrow{\pi}_q$ determined kinetically in collisional equilibrium ($\partial/\partial t \ll \nu$).

Two-Fluid Moment Equations For Extended MHD

- Exact fluid moment equations for each plasma species result from velocity moments ($\int d^3v \vec{v}^n$, $n = 0, 1, 2$) of the plasma kinetic equation:

$$\begin{aligned}
 n = 0, |\vec{v}|^0, \text{ density} \quad & \frac{\partial n}{\partial t} + \vec{\nabla} \cdot n \vec{V} = 0, \quad \{\vec{\nabla} \vec{V}\} \equiv \frac{1}{2} \left[\vec{\nabla} \vec{V} + (\vec{\nabla} \vec{V})^T \right] - \frac{1}{3} \overleftrightarrow{\mathbf{I}} (\vec{\nabla} \cdot \vec{V}), \\
 n = 1, \vec{v}, \text{ momentum} \quad & mn \frac{d\vec{V}}{dt} = nq(\vec{E} + \vec{V} \times \vec{B}) - \vec{\nabla} p - \vec{\nabla} \cdot \pi + \vec{R}, \quad \frac{d}{dt} \equiv \frac{\partial}{\partial t} + \vec{V} \cdot \vec{\nabla}, \\
 n = 2, v^2, \text{ energy} \quad & \frac{3}{2} n \frac{dT}{dt} + nT \vec{\nabla} \cdot \vec{V} = -\vec{\nabla} \cdot \vec{q} - \overleftrightarrow{\pi} : \{\vec{\nabla} \vec{V}\} + Q, \quad p \equiv nT, \\
 \text{or entropy} \quad & \frac{\partial(ns)}{\partial t} + \vec{\nabla} \cdot \left(ns \vec{V} + \frac{\vec{q}}{T} \right) = \frac{1}{T} (-\vec{q} \cdot \vec{\nabla} \ln T - \overleftrightarrow{\pi} : \{\vec{\nabla} \vec{V}\} + Q), \quad s \equiv \ln(T^{3/2}/n).
 \end{aligned}$$

- Collisional friction forces and energy exchange terms including only flow, heat flow effects (i.e., neglecting energy-weighted heat flow etc.) are:

$$\begin{aligned}
 \vec{R}_e &= -m_e n_e \nu_e \left[(\vec{V}_e - \vec{V}_i) - \frac{3}{5 n_e T_e} \vec{q}_e \right] = \frac{n_e e}{\sigma_0} \left(\vec{J} + \frac{3e}{5T_e} \vec{q}_e \right), \quad \sigma_0 \equiv \frac{n_e e^2}{m_e \nu_e}, \quad \vec{R}_i = \vec{R}_e, \\
 Q_e &= -Q_i - (\vec{V}_e - \vec{V}_i) \cdot \vec{R}_e = -Q_i + \frac{1}{\sigma_0} \left(|\vec{J}|^2 + \frac{3e}{5T_e} \vec{J} \cdot \vec{q}_e \right), \quad Q_i = \frac{3}{2} n_e \nu_e (T_e - T_i).
 \end{aligned}$$

- The moment equations need closure moments for \vec{q} and $\overleftrightarrow{\pi}$ ($\vec{v}_r \equiv \vec{v} - \vec{V}$):

$$\text{heat flux } \vec{q} \equiv \int d^3v \vec{v}_r \left(\frac{m v_r^2}{2} - \frac{5}{2} \right) f, \quad \text{stress tensor } \overleftrightarrow{\pi} \equiv \int d^3v m \left(\vec{v}_r \vec{v}_r - \frac{v_r^2}{3} \overleftrightarrow{\mathbf{I}} \right) f.$$

Neoclassical Closures

- There are two basic approaches to neoclassical transport theory for axisymmetric toroidal plasmas:

kinetic [9] — collisions of particles drifting off flux surfaces cause radial transport,
fluid [10] — viscous drag on untrapped particles due to collisions with “immobile”
 trapped particles causes parallel/poloidal force that leads to radial plasma transport.

- Only fluid moment approach is relevant for Extended MHD equations.
- Key assumptions in deriving usual neoclassical closures for transport:
 axisymmetric magnetic field geometry — no ripples, $\delta\vec{B}$, or magnetic islands,
 collisional equilibrium ($\partial/\partial t < \nu$) between trapped and (flowing) untrapped particles,
 flux-surface-average is appropriate because on collisional time scale the particles
 circumnavigate poloidal cross-section of torus many times — $\lambda \equiv v_T/\nu \gg 2\pi R_0 q$.
- The relevant neoclassical closures are the flux-surface-average of the parallel viscous forces induced by the poloidal flow U_θ and heat flow Q_θ :

$$\begin{pmatrix} \langle \vec{B} \cdot \vec{\nabla} \cdot \vec{\pi}_{\parallel} \rangle \\ \langle \vec{B} \cdot \vec{\nabla} \cdot \vec{\Theta}_{\parallel} \rangle \end{pmatrix} = mn \langle B^2 \rangle \begin{pmatrix} \mu_{00} & \mu_{01} \\ \mu_{10} & \mu_{11} \end{pmatrix} \begin{pmatrix} U_\theta \\ Q_\theta \end{pmatrix}, \quad \mu \sim \sqrt{\epsilon} \nu, \quad U_\theta \equiv \frac{\vec{V} \cdot \vec{\nabla} \theta}{\vec{B} \cdot \vec{\nabla} \theta} = \frac{V_{\parallel}}{B} + \frac{\vec{V}_{\perp} \cdot \vec{\nabla} \theta}{\vec{B} \cdot \vec{\nabla} \theta}.$$

[9] F.L. Hinton and R.D. Hazeltine, Rev. Mod. Phys. 48, 239 (1976) — kinetic neoclassical transport.

[10] S.P. Hirshman and D.J. Sigmar, Nuclear Fusion 21, 1079 (1981) — fluid moment neoclassical transport.

Neoclassical MHD Is One Set Of Extended MHD Equations

- Neoclassical MHD equations [11] use collisionally equilibrated parallel viscous forces and approximate gyroviscous forces to yield (for $t > 1/\nu$):
 neoclassical parallel Ohm's law, including trapped particle effects on electrical conductivity [$\sigma \simeq \sigma_{\text{Sp}}/(1 + \mu_e/\nu_e)$] and bootstrap current [$J_{\text{bs}} \sim (\mu_e/\nu_e)dP/d\psi$], poloidal flow damped to zero [or small $V_\theta \sim (1.17/q_i)dT_i/d\psi$] \implies only toroidal flow, enhanced \perp dielectric and inertia $\implies 1 + c^2/c_A^2 \implies 1 + c^2/c_{A\theta}^2$, larger by $B^2/B_\theta^2 \sim 10^2$, neoclassical tearing modes (NTMs) driven by island perturbation of bootstrap current.

- Hegna suggested a heuristic local (i.e., not flux-surface-averaged) neoclassical closure to facilitate NIMROD simulations of NTMs [12]:

$$\vec{\nabla} \cdot \vec{\pi}_{\parallel} = mn\mu \langle B^2 \rangle \frac{\vec{V} \cdot \vec{e}_\theta}{(\vec{B} \cdot \vec{e}_\theta)^2} \vec{e}_\theta, \quad \vec{e}_\theta \equiv \sqrt{g} \frac{\vec{\nabla} \zeta \times \vec{\nabla} \psi}{\vec{B} \cdot \vec{\nabla} \theta}.$$

- To proceed further with neoclassical MHD-type simulations we need:
 time-dependent parallel viscous force — to study NTM threshold behavior,
 local parallel viscous force (or pressure-anisotropy) — to facilitate local analysis,
 inclusion of nonaxisymmetric effects of islands, $\delta \vec{B}$ — for toroidal flow damping.

[11] J.D. Callen, W.X. Qu, K.D. Siebert, B.A. Carreras, K.C. Shaing and D.A. Spong in *Plasma Physics and Controlled Nuclear Fusion Research 1986* (IAEA, Vienna, 1987), Vol. 2, p 157.

[12] T.A. Gianakon, S.E. Kruger and C.C. Hegna, *Phys. Plasmas* 9, 536 (2002) — NIMROD sim. of NTMs.

Recent Progress On Neoclassical Closures

- Temporal behavior of the parallel viscous force has been calculated [13], with the most useful formulas obtained in a small $\epsilon \equiv \Delta B/2B$ expansion:

$$\langle \vec{B} \cdot \vec{\nabla} \cdot \overleftrightarrow{\pi} \rangle \simeq mn \langle B^2 \rangle \mu \left[U_\theta(t) + \frac{1}{\bar{\nu}} \frac{\partial U_\theta}{\partial t} + \sum_n c_n \int_0^t d\tau e^{-\bar{\nu} \kappa_n (t-\tau)} \frac{\partial U_\theta}{\partial t} \right],$$

which implies a “time-history” integral equation for the parallel flow evolution,

$$U_\theta(t) = h_\theta(t) + \int_0^t d\tau K_\theta(t; \tau) U_\theta(\tau) \text{ in which } h_\theta \text{ represents initial conditions.}$$

- Local pressure anisotropy $\pi_{\parallel} \equiv p_{\parallel} - p_{\perp}$ also determined recently [14]:

$\pi_{\parallel} \sim mn\mu U_\theta \langle B^2 \rangle \times$ (incomplete elliptic functions) — continuous function of θ ;

however, $\vec{B} \cdot \vec{\nabla} \cdot \overleftrightarrow{\pi}_{\parallel} = (2/3) \vec{B} \cdot \vec{\nabla} \pi_{\parallel} + \pi_{\parallel} (\vec{B} \cdot \vec{\nabla} \ln B)$ is divergent at $B = B_{\max}$.

- Nonaxisymmetry effects introduced by magnetic islands or $\delta \vec{B}$ cause neoclassical radial particle fluxes, toroidal viscous flow damping [15,16]:

Magnetic islands change radial location of $\psi \implies \partial(\vec{V} \cdot \vec{\nabla} \zeta) / \partial t \sim (w/a)^2 \sim (\delta B_{x_{mn}}/B)$;

Ideal MHD $\delta \vec{B}$ produces helical change in equilibrium $\implies \partial(\vec{V} \cdot \vec{\nabla} \zeta) / \partial t \sim |\delta \vec{B}/B|^2$.

[13] A.L. Garcia-Perciante, J.D. Callen, K.C. Shaing, and C.C. Hegna, Phys. Plasmas 12, 052516 (2005).

[14] A.L. Garcia-Perciante, J.D. Callen, K.C. Shaing, and C.C. Hegna, Phys. Plasmas 13, 012509 (2006).

[15] K.C. Shaing, Phys. Rev. Lett. 87, 245003 (2001) — particle flux due to magnetic islands.

[16] K.C. Shaing, Phys. Plasmas 10, 1443 (2003) — toroidal momentum damping due to MHD modes.

Complete Set Of Extended MHD (ExMHD) Equations

- Combining plasma fluid and Maxwell's equations, one obtains the complete set of "Extended MHD" equations:

density $\frac{\partial \rho_m}{\partial t} + \vec{\nabla} \cdot \rho_m \vec{V} = 0,$

momentum $\rho_m \frac{d\vec{V}}{dt} = \vec{J} \times \vec{B} - \vec{\nabla} P - \vec{\nabla} \cdot \overleftrightarrow{\Pi}, \quad P \equiv p_e + p_i, \quad \overleftrightarrow{\Pi} \simeq \overleftrightarrow{\pi}_i + \overleftrightarrow{\pi}_e \simeq \overleftrightarrow{\pi}_i,$

magnetic field $\frac{\partial \vec{B}}{\partial t} = -\vec{\nabla} \times \vec{E}, \quad \vec{J} = \vec{\nabla} \times \vec{B} / \mu_0, \quad \vec{\nabla} \cdot \vec{B} = 0,$

Ohm's law $\vec{E} = -\vec{V} \times \vec{B} + \frac{\vec{R}_e}{n_e e} + \frac{\vec{J} \times \vec{B} - \vec{\nabla} p_e - \vec{\nabla} \cdot \overleftrightarrow{\pi}_e}{n_e e} + \frac{m_e}{e^2} \frac{d}{dt} \left(\frac{\vec{J}}{n_e} \right).$

Eq. of state $\frac{d}{dt} \left(\ln \frac{P}{\rho_m^\Gamma} \right) = \frac{\Gamma - 1}{P} \left(p_e \frac{ds_e}{dt} + p_i \frac{ds_i}{dt} \right), \quad \Gamma = \frac{5}{3}.$

Additional Specifications Needed For ExMHD Equations

- Electron temperature T_e , pressure $p_e = n_e T_e$, flow $\vec{V}_e \equiv -\vec{J}/n_e e + \vec{V}_i$:

$$\frac{dT_e}{dt} \equiv \frac{\partial T_e}{\partial t} + \vec{V}_e \cdot \vec{\nabla} T_e = \frac{2}{3} T_e \left(-\vec{\nabla} \cdot \vec{V}_e + \frac{ds_e}{dt} \right) \implies \frac{3}{2} \frac{\partial p_e}{\partial t} = -\vec{V}_e \cdot \vec{\nabla} p_e - \vec{\nabla} \cdot \left(\frac{5}{2} p_e \vec{V}_e \right) + p_e \frac{ds_e}{dt}$$

- Electron entropy $s_e \equiv \ln(T_e^{3/2}/n_e)$:

$$\frac{ds_e}{dt} \equiv \frac{\partial s_e}{\partial t} + \vec{V}_e \cdot \vec{\nabla} s_e = -(\vec{\nabla} \cdot \vec{q}_e + \vec{\pi}_e : \vec{\nabla} \vec{V}_e - Q_e)/n_e T_e$$

- Ion temperature T_i , pressure $p_i = n_i T_i$, flow $\vec{V}_i \simeq \vec{V}$:

$$\frac{dT_i}{dt} \equiv \frac{\partial T_i}{\partial t} + \vec{V}_i \cdot \vec{\nabla} T_i = \frac{2}{3} T_i \left(-\vec{\nabla} \cdot \vec{V}_i + \frac{ds_i}{dt} \right) \implies \frac{3}{2} \frac{\partial p_i}{\partial t} = -\vec{V}_i \cdot \vec{\nabla} p_i - \vec{\nabla} \cdot \left(\frac{5}{2} p_i \vec{V}_i \right) + p_i \frac{ds_i}{dt}$$

- Ion entropy $s_i \equiv \ln(T_i^{3/2}/n_i)$:

$$\frac{ds_i}{dt} \equiv \frac{\partial s_i}{\partial t} + \vec{V}_i \cdot \vec{\nabla} s_i = -(\vec{\nabla} \cdot \vec{q}_i + \vec{\pi}_i : \vec{\nabla} \vec{V}_i - Q_i)/n_i T_i$$

- Collisional friction force \vec{R}_e :

$$\vec{R}_e = \frac{n_e e}{\sigma_0} \left(\vec{J} + \frac{3e}{5T_e} \vec{q}_e \right), \quad \sigma_0 \equiv \frac{n_e e^2}{m_e \nu_e}$$

- Collisional energy exchange:

$$Q_e = -Q_i + \frac{1}{\sigma_0} \left(|\vec{J}|^2 + \frac{3e}{5T_e} \vec{J} \cdot \vec{q}_e \right), \quad Q_i = \frac{3}{2} n_e \nu_e (T_e - T_i).$$

Extended MHD Equations Require Various Closures

- Closures for \vec{q} and $\overleftrightarrow{\pi}$ need to have their parallel (\parallel), cross (\wedge) and perpendicular (\perp) components specified:

$$\vec{q} = \vec{q}_{\parallel} + \vec{q}_{\wedge} + \vec{q}_{\perp}, \quad \text{and} \quad \overleftrightarrow{\pi} = \overleftrightarrow{\pi}_{\parallel} + \overleftrightarrow{\pi}_{\wedge} + \overleftrightarrow{\pi}_{\perp}.$$

- Parallel heat flow $\vec{q}_{\parallel} = q_{\parallel} \vec{b}$, $\vec{b} \equiv \vec{B}/B$, $q_{\parallel} \equiv -\int d^3v v_{\parallel} L_1^{3/2} F$ determined using F obtained solving a Chapman-Enskog-type drift kinetic equation [4,5]:

$$\frac{\partial F}{\partial t} + v_{\parallel} \vec{b} \cdot \vec{\nabla} F = \mathcal{C}_R\{F\} + v_{\parallel} L_1^{(3/2)} f_M \vec{b} \cdot \vec{\nabla} T - \frac{m}{T} \left(v_{\parallel}^2 - \frac{v_{\perp}^2}{2} \right) f_M (\vec{b} \cdot \{\vec{\nabla} \vec{V}\} \cdot \vec{b}) + \dots$$

- Various approaches used to obtain q_{\parallel} from this parallel kinetic equation:

Collisional regime (Braginskii) — neglect $\partial F/\partial t$, $v_{\parallel} \vec{b} \cdot \vec{\nabla} F$; invert collision operator;

Collisionless — linearize and obtain Hammett-Perkins [17] Landau-type closures [5,18];

PIC-type δf code (Barnes) — but higher order moments are “noisier?”

“Continuum” type solutions [19] — expand F in pitch-angle eigenfunctions of $\mathcal{C}_R\{F\}$.

- Stress $\overleftrightarrow{\pi}_{\parallel} \equiv \pi_{\parallel} (\vec{b}\vec{b} - \mathbf{I}/3)$, $\vec{b} \equiv \vec{B}/B$, $\pi_{\parallel} \equiv p_{\parallel} - p_{\perp} = \int d^3v m (v_{\parallel}^2 - v_{\perp}^2/2) F$ is also determined from the solution of the parallel kinetic equation [20] \implies neoclassical closures for π_{\parallel} and $\langle \vec{B} \cdot \vec{\nabla} \cdot \overleftrightarrow{\pi}_{\parallel} \rangle$ for $\omega \ll \nu \sim k_{\parallel} v_T$.

- [17] G.W. Hammett and F.W. Perkins, Phys. Rev.Lett. 64, 3019 (1990) — simplest Landau closure for q_{\parallel} .
 [18] A.I. Smolyakov, M. Yagi, J.D. Callen, Fields Inst. Comm. 46, 243 (2005) — neutral fluid nonlocal clos.
 [19] E.D. Held *et al.*, Phys. Plasmas 11, 2419 (2004) and references cited therein — q_{\parallel} closure, general $\lambda \nabla_{\parallel}$.
 [20] E.D. Held, “Unified form for parallel ion viscous stress in magnetized plasmas,” PoP 10, 4708 (2003).

Some Complications In Obtaining Parallel Closures

- The “usual” Chapman-Enskog-like drift-kinetic equation (DKE) [4,5] has many ($\gtrsim 5$) “drives” on its right side in terms of the form

$v_{\parallel} f_M \vec{B} \cdot \vec{\nabla} \cdot \overleftrightarrow{\pi} \implies$ causes $\sim \sqrt{\epsilon}$ correction to parallel viscous forces,

$v_{\parallel} R_{e\parallel} \implies$ additional corrections to parallel flow?, part of Spitzer problem?

dissipative $L_1^{(1/2)}$ terms due to $\overleftrightarrow{\pi} : \{ \vec{\nabla} \vec{V} \}$, $\vec{\nabla} \cdot \vec{q}$ and $Q \implies$ temperature change δT ?

- Also, ϵ_{\perp}^2 additions to DKE — Catto & Simakov [21], paleoclassical [22].
- Unfortunately, to obtain parallel closures to sufficient accuracy ($\sim \tilde{\epsilon}_{\perp}^2 \tilde{\epsilon}_{\parallel} n \sim \epsilon_{\perp}^2$) one needs to keep $\mathcal{O}(\tilde{\epsilon}_{\perp}^2)$ terms, particularly for $n \gg 1$.
- Shaing and Spong [3] have exhibited some of the 3D complications that arise in long collision length plasmas by obtaining a “local” closure relation for $\pi_{\parallel} \equiv p_{\parallel} - p_{\perp}$ in the plateau collisionality regime.
- Also, Shaing emphasizes that in general there are not enough free parameters in kinetic analysis to satisfy all Chapman-Enskog constraints \implies residual “extra” (but usually higher order) δn , δT , $\delta \vec{V}$ terms.

[21] A.N. Simakov, P.J. Catto, Phys. Plasmas 12, 012105 (2005) — additional $\mathcal{O}(\epsilon_{\perp}^2)$ terms in DKE (for $\overleftrightarrow{\pi}_{\wedge}$).

[22] J.D. Callen, Phys. Rev. Lett. 94, 055002 (2005); Nucl. Fus. 45, 1120 (2005); Phys. Pl. 12, 092512 (2005).

Extended MHD Equations Require Various Closures (cont'd)

- Components of \vec{q} perpendicular to \vec{B} can be obtained from the fluid moment equation for $\partial\vec{q}/\partial t$ [4], $\vec{R}_q \sim mn\nu[l_{10}(\vec{V}_e - \vec{V}_i) + l_{11}(-2\vec{q}_e/5p_e)]$:

$$\frac{d\vec{q}}{dt} = \frac{\omega_c}{B}\vec{q} \times \vec{B} - \frac{5nT}{2m}\vec{\nabla}T + \frac{T}{m}(\vec{\nabla} \cdot \overleftrightarrow{\Theta} + \vec{R}_q) - \dots, \text{ which upon taking } \vec{B} \times \text{ yields}$$

$$\vec{q}_\wedge = \underbrace{\frac{5\vec{B} \times \vec{\nabla}T}{2\omega_c}}_{\text{diamagnetic}} \sim \epsilon_\perp, \quad \text{and} \quad \vec{q}_\perp = \frac{1}{\omega_c B} \vec{B} \times \left[\frac{T}{m} \underbrace{(\vec{R}_q + \vec{\nabla} \cdot \overleftrightarrow{\Theta})}_{\text{classical + neo}} - \frac{d\vec{q}}{dt} + \dots \right] \sim \epsilon_\perp^2.$$

- A similar analysis of the $d\overleftrightarrow{\pi}/dt$ equation can be performed to yield

$$\overleftrightarrow{\pi}_\wedge = \underbrace{\frac{2p}{\omega_c} \overleftrightarrow{\mathcal{K}}^{-1} \{ \{ \vec{\nabla} \vec{V} \} + \frac{4}{5nT} \{ \vec{\nabla} \vec{q} \} }}_{\text{gyroviscous stress}} \sim \epsilon_\perp, \quad \text{and} \quad \overleftrightarrow{\pi}_\perp = \underbrace{\frac{\nu_{\text{eff}}}{\omega_c} \overleftrightarrow{\mathcal{K}}^{-1} \{ \overleftrightarrow{\pi}_\wedge \} + \dots}_{\text{perpendicular stress}} \sim \epsilon_\perp^2,$$

in which the inverse tensor operator [23] $\overleftrightarrow{\mathcal{K}}^{-1} \{ \overleftrightarrow{\mathbf{S}} \} = \frac{1}{4} \left([\vec{b} \times \overleftrightarrow{\mathbf{S}} \cdot (\vec{\mathbf{I}} + 3\vec{b}\vec{b})] + \text{transpose} \right)$.

- Recently, Ramos [24] used fluid moments to obtain a compact form for the gyroviscous stress tensor $\overleftrightarrow{\pi}_\wedge$ for arbitrary magnetic geometry.

[23] C.T. Hsu, R.D. Hazeltine and P.J. Morrison, Phys. Fluids 29, 1480 (1986) — see Appendix A.

[24] J.J. Ramos, “Fluid formalism for collisionless magnetized plasmas,” Phys. Plasmas 12, 052102 (2005).

Inclusion of Micro-turbulence Effects in ExMHD?

- Can effects of micro-turbulence be included in ExMHD simulations?

Neutral fluid turbulence closure models seek Reynolds stress closures that represent non-dissipative transfer of energy to higher \vec{k} in inertial range, then dissipation.

Drift-wave-type micro-turbulence has unstable \vec{k} -space region ($k_{\perp} \rho_S \sim 0.2-1$) but some mode coupling (inverse cascade) to other \vec{k} (e.g., to $k_{\theta} = k_{\zeta} = 0$ zonal flows) — can these reactive, inverse cascade and dissipative effects be included in ExMHD?

- Some ExMHD physical effects expected from micro-turbulence:

Flow damping (or growth! [25]) induced by Reynolds stress

Radial particle, electron and ion heat, and momentum diffusion via anomalous D_a , χ_e and χ_i , and χ_{ϕ}

- By averaging over microscopic scales [25]? — still being worked on:

Separate [25] wavelength ranges of density via $n = n^{<} + n^{>}$ with $n^{<}$ including all \vec{k} up to say $k_{\theta} \rho_S \leq 0.2$ and $n^{>}$ representing all higher \vec{k} processes (i.e., micro-turbulence).

Then, average over high \vec{k} to obtain effects of micro-turbulence in the macroscopic description: $\partial n^{<}/\partial t + \vec{\nabla}^{<} \cdot n^{<} \vec{V}^{<} + \vec{\nabla}^{<} \cdot \vec{\Gamma}_a = 0$ in which $\vec{\Gamma}_a = \langle n^{>} \vec{V}^{>} \rangle_{>} \simeq -D_a \vec{\nabla}^{<} n^{<}$.

Similarly, Reynolds stress and heat flux terms are added to momentum, energy eqs.

[25] C.J. McDevitt and P.H. Diamond, “Multi-Scale Interaction of a Tearing Mode with Drift Wave Turbulence: A Minimal Self-Consistent Model,” *Phys. Plasmas* 13, 032302 (2006).

There Are Two Generic Types Of Extended MHD Problems

- “Fast MHD” ($\omega > \nu_i \sim 10^3 \text{ s}^{-1}$) phenomena occur on Alfvénic timescale:

Examples: sawtooth crashes, disruption precursors (DIII-D #87009), ELMs.

Physically, need ideal MHD plus diamagnetic flow & gyroviscosity (two-fluid) effects — ω_{*i} stabilization for 1/1 sawtooth crashes, plus ω_{*e} for stabilizing high mode numbers.

Dissipative closures operate on longer time scales ($t > 1/\nu$) and hence are negligible — except for parallel T_e equilibration in irregular magnetic fields, destabilizing resistivity effects, and possibly stabilizing diffusive effects on high mode numbers [26].

- “Slow MHD” ($\omega < \nu_i \sim 10^3 \text{ s}^{-1}$) phenomena occur on the resistive time scale and involve many physical processes:

Examples: $\Delta' > 0$ tearing modes, NTMs, RWMs.

Neoclassical MHD effects important — poloidal flow damping \implies only toroidal flow, enhanced inertia (by $B^2/B_\theta^2 \gg 1$), neoclassical parallel resistivity, bootstrap current.

Since nonlinear evolution (tearing modes \implies magnetic islands, RWMs \implies kink in plasma growing on resistive wall time $\sim 10 \text{ ms}$) is on transport time scale, all transport effects are important — need complete (\parallel, \wedge, \perp) dissipative closures for $\vec{q}, \vec{\pi}$.

Diamagnetic flow (ω_*) effects are ultimately not so critical — vanish on separatrix where $\vec{\nabla}P \rightarrow 0$, or just lead to slight changes in toroidal flow velocity.

The second order (in gyroradius) effects are needed for perturbations, but they may not be needed for equilibrium since they represent negligible classical diffusion effects.

[26] B.A. Carreras, L. Garcia and P.H. Diamond, Phys. Fl. 30, 1388 (1987) — χ_\perp, μ_\perp effects on res-g modes.

Discussion: Develop Closures For Classes Of Problems?

- Closures can only be systematically derived for collisional regime \implies Braginskii equations — but toroidal plasmas violate $\lambda \nabla_{\parallel} < 1$ condition.
- No general closures can be derived for long collision length λ regimes — because \parallel kinetics depends on geometry over the collision length.
- Also, needed (for \wedge , \perp closures) first and second order terms in finite gyroradius expansion are complicated and depend on gradients of \vec{B} .
- Thus, one is led to consider key closures needed for classes of problems:
 - Fast MHD ($\omega > \nu_i$)
mainly just diamagnetic flows and lowest order gyroviscosity, but perhaps with some diffusivities to stabilize high mode numbers.
 - Slow MHD ($\omega < \nu_i$)
tearing modes, NTMs — mainly just neo \parallel viscous force (but local with dynamics),
RWMs — mainly just equilibrium neoclassical parallel viscous force, plus toroidal flow damping induced as mode kinks the plasma and magnetic field.
- But for ultimate extended MHD simulations of toroidal plasmas one will need to develop procedures for determining and numerically implementing \parallel , \wedge and \perp closures (& extra terms) to sufficient accuracy.

Closures For NIMROD Have Special Forms, Requirements

- They should be developed with “drives” due to $\vec{\nabla}n$, $\vec{\nabla}T$, $\vec{\nabla}\vec{V}$ — i.e., via a Chapman-Enskog-type approach (at least for parallel kinetics).
- They should be written in real space \vec{x} and time t — i.e., not in \vec{k}, ω .
- The various parts of the closures should be handled differently:
 - parallel: C-E type drift-kinetic analysis must be used, including collisional effects,
 - cross: fluid-type analysis, gyroviscosity for these diamagnetic flow type effects,
 - perpendicular: fluidlike radial transport due to collisional effects on diamagnetic flows, perhaps plus anomalous transport fluxes due to micro-turbulence.
- There is no need to determine divergences of the closures (i.e., $\vec{\nabla} \cdot \vec{q}$ and $\vec{\nabla} \cdot \overleftrightarrow{\pi}$), since for finite element representations (ξ) such as in NIMROD, an integration by parts yields integrals that only depend on \vec{q} and $\overleftrightarrow{\pi}$:
$$\int d^3x \xi \vec{\nabla} \cdot \vec{q} = \int_V d^3x \vec{q} \cdot \vec{\nabla} \xi + \iint_S d\vec{S} \cdot \xi \vec{q},$$
 and similarly for $\vec{\nabla} \cdot \overleftrightarrow{\pi}$.
- Extended MHD equations with closures should clearly satisfy momentum and energy conservation relations for the overall system — plasma plus electromagnetic fields

Summary

- Extended MHD model from two-fluid equations with $\vec{R}_e = \frac{n_e e}{\sigma_0} (\vec{J} + \frac{3e}{5T_e} \vec{q}_e)$ provide basis for describing macroscopic plasma behavior — if suitable closures for \vec{q} and $\vec{\pi}$ are available and/or numerically implementable.
- Rigorous analysis in collisional regime (Braginskii) shows that closures in a magnetized plasma are very anisotropic — $\vec{q}_{\parallel} \sim \epsilon_{\perp}^0 \epsilon_{\parallel}^2$ (parallel), $\vec{q}_{\wedge} \sim \epsilon_{\perp}$ (diamagnetic), $\vec{q}_{\perp} \sim \epsilon_{\perp}^2$ (perpendicular) and similarly for $\vec{\pi}$.
- Determinations of closure components depend on direction:
 - || (kinetic, $\epsilon_{\parallel} \sim |\lambda \nabla_{\parallel}| \gtrsim 1$) — parallel Chapman-Enskog-type drift-kinetic equation,
 - \wedge (diamagnetic, $\epsilon_{\perp} \sim |\rho \vec{\nabla}_{\perp}| \ll 1$) — can use $\vec{B} \times$ fluid moment equations for \vec{q} , $\vec{\pi}$,
 - \perp (perp, $\epsilon_{\perp}^2 \ll 1$) — gyroradius smaller parts of $\vec{B} \times$ same fluid moments.
- A comprehensive set of Extended MHD (ExMHD) equations, including micro-turbulence effects, are being developed — but numerical procedures for determining parallel closures are needed for completeness.
- ExMHD equations, closures for NIMROD have special requirements.