


Theme: Codes Verification/Validation Should be a Collective Effort

N. N. Gorelenkov for CSEPP


Princeton Plasma Physics Laboratory, Princeton

CSEPP kickoff meeting, Boulder, CO

March 29, 2008



General scheme for problem oriented V&V



- identify problem(s)
 - identify properties and requirements for the model,
 - apply codes.
- There are some missing links:
 - linear global codes with FLR effects,
 - multiple AE instability codes.
- GKM code is the ultimate goal
⇒ should be kept in mind while developing the plasma scenarios.

Proper plasma conditions are required on initial runs.
Information should be shared (project website?).

What do we compare?

Synthetic diagnostic is required for V&V: such as for comparison with plasma diagnostics - should not be overlooked.

On early stage of code development.

What do we compare?

Synthetic diagnostic is required for V&V: such as for comparison with plasma diagnostics - should not be overlooked.

On early stage of code development.

- Instabilities/modes we see in codes has to be identified based on mode properties:
 - frequency
 - structure
 - polarization
 - plasma parametric dependencies

What do we compare?

Synthetic diagnostic is required for V&V: such as for comparison with plasma diagnostics - should not be overlooked.

On early stage of code development.

- Instabilities/modes we see in codes has to be identified based on mode properties:
 - frequency
 - structure
 - polarization
 - plasma parametric dependencies
- Nonlinear benchmarks, validations are challenging, need cross (nonlinear) code comparisons - fewer examples exist, less agreement?
 - saturation of modes: single to multiple,
 - DIII-D, NSTX transport problems,
 - predictive simulation is the goal.

Need to prioritize potential problems:

Need to prioritize potential problems:

*AEs, potentially having high priority

- TAEs, EAEs, NAEs...
 - low-n modes or medium-n low shear: simple 2 harmonic coupling.

Need to prioritize potential problems:

*AEs, potentially having high priority

- TAEs, EAEs, NAEs...
 - low- n modes or medium- n low shear: simple 2 harmonic coupling.
- RSAEs/Alfven cascades
 - dominant single m , analytically treatable including kinetic physics
 - * (potentially more) important for EP transport.

Need to prioritize potential problems:

*AEs, potentially having high priority

- TAEs, EAEs, NAEs...
 - low- n modes or medium- n low shear: simple 2 harmonic coupling.
- RSAEs/Alfven cascades
 - dominant single m , analytically treatable including kinetic physics
 - * (potentially more) important for EP transport.
- BAE/BAAEs
 - coupling to acoustic wave, not well understood.
 - although may not be expected in ITER.

Candidate problems (part 2)

- $m = 1$ kink modes (IFS lead) - ITER is interested:
 - sawtooth, fishbones,
 - a lot of studies, existing benchmarks (Porcelli, Mcclements, NOVA-KN, M3D)
 - complicated nonlinear dynamics (Odblom),
 - acoustic mode coupling: γ is known to effect the growth rate of $m = 1$ kink drive and threshold.

Candidate problems (part 2)

- $m = 1$ kink modes (IFS lead) - ITER is interested:
 - sawtooth, fishbones,
 - a lot of studies, existing benchmarks (Porcelli, Mcclements, NOVA-KN, M3D)
 - complicated nonlinear dynamics (Odblom),
 - acoustic mode coupling: γ is known to effect the growth rate of $m = 1$ kink drive and threshold.
- Plasma turbulence vs alphas.
- Ion Cyclotron Emission - cyclotron range of frequency.
- Integrated modeling.
- α -channeling.

First year work

- Large aspect ratio circular plasma, analytical pressure, q , density profiles → to realistic plasma (see later)
 - TAEs, RSAEs (*Gorelenkov, PPCF'06*)
 - * Linear theory is applicable in ideal MHD limit plus perturbative kinetic effects.
 - * Linear codes are applicable. FLR can be included in some codes, LIGKA may be involved.
 - * Nonlinear theory can be applied for single mode saturation via NOVA evaluation of theoretical results. ORBIT also can be used for cross benchmarks.
 - $m = 1$ mode (IFS), NOVA(-KN) to be applied.

Further work

Alfven - acoustic coupling, effect on $m = 1$, low - f modes.

Validation against experiments:

- TAE, RSAE in DIII-D, TFTR, NSTX ... for saturation BAAE/BAE

Further work

Some missing links:


- kinetic global non-perturbative code for high-, medium- n modes,
- can modify NOVA and use its infrastructure,
- two fluid approach should give Pade like kinetic effects.

Some examples


*Make use of present documented cases (M. Van Zeeland)
MHD TAE structure seems to agree with measurements in DIII-D*

Internal TAE/RSAEs mode structures measured by ECE, compared with NOVA

RSAE & TAE frequencies crossover




RSAE & TAE structures



Saturation is measured, single, multiple mode saturation can be obtained.
Transport is not explained.

ORBIT comparison with quasilinear theory (IFS) using NOVA-K

PPPL



- Single mode saturation via EP response can be compared with theory.
- Improved NOVA-K simulation could include mapping techniques.
- Can be used for validation with single mode XPs.

*Some modeling challenges for *AE instabilities*

- Beams, ICRH ions are often superalfvenic. How does their distribution evolve?
- How well do we know (XP & theory) the mode structure? (Carolipio'01, Heidbrink'97)
- What about thermal ions? Will they drive modes as in DIII-D experiments (Nazikian'06)?
- Non-ideal damping mechanism can be addressed in MHD codes only perturbatively.
- Damping models need to be validated against experiments for medium to high toroidal, n , numbers.
- High- n ideal MHD codes need high poloidal harmonic number up to $m = 100$ or high poloidal grid resolution.

*Some modeling challenges for *AE instabilities*

NOVA/NOVA-K is hybrid kinetic/MHD code with perturbative treatment of various kinetic effects.

Its development is based on **multi-institutional collaborations** and includes experimental validations:



- NOVA uses **TRANSP** output for plasma parameters
- Mode structure is computed within ideal MHD (NOVA)
- Perturbative kinetic mode analysis is performed with NOVA-K code
- Fast ion drive includes: finite orbit width (FOW) and FLR effects
- Damping mechanisms included are
 - ion/electron Landau
 - radiative, non-ideal
 - trapped electron collisional
 - continuum damping
- Nonlinear saturation quasilinear model is available...

Typical experimental conditions: multiple modes are present

PPPL

DIII-D/NSTX similarity experiments were designed to study machine size scaling predictions (W. Heidbrink, PPCF '03):

- The same minor radius in NSTX and DIII-D but different major radii.
- Use similar NBI features: injection geometry, energy, trapped to passing particle ratio.



Most unstable mode number (larger amplitude at the edge) scales as $n \sim a/q^2$.



How do we model EP distribution function (df)

PPPL

- Alphas in burning plasma (BP) will have slowing down df.
- ICRH df is an issue.
- NBI EP df model exists, should be used.

*Truncated image method
vs. Monte-Carlo*

*TRANSP Lorentz diffusion operator
is consistent with image method (ITER NBI)*




Great reduction of Monte-Carlo noises in TRANSP simulations. Analytical derivatives can be taken.

DIII-D recent RS plasma XP (W.W. Heidbrink, IAEA'06)


PPPL

The Fast-ion Density Gradient is Flattened



- The profile remains flat during the strongest Alfvén activity

- As the activity weakens the profile peaks but is still broader than classically predicted




**For this comparison, the FIDA density profile is normalized to the equilibrium profile at 1.20 s.*

NSTX multi-mode transport XPs

PPPL


Multi-mode driven transport is targeted on NSTX M3D study is in progress



NSTX multi-mode transport XPs

PPPL

Multi-mode driven transport is targeted on NSTX M3D study is in progress



At the moment the diagnostic has to be advanced for better internal measurements.

NSTX is committed for EP effects studies: multi-mode effects on EP transport and current drive.

Summary of what NOVA-K can do

PPPL

- Linear theory with various driving and damping mechanisms of *AEs (1st year).
- Employs quasilinear theory (IFS) for TAE saturation (2nd-3rd year).
- Relatively fast, easy to modify.