

Velocity-space compression from Fermi acceleration with Lorentz scatteringJ. C. Waybright, M. E. Mlodik, and N. J. Fisch *Department of Astrophysical Sciences, Princeton University, Princeton, New Jersey 08540, USA*

(Received 3 September 2021; accepted 5 January 2022; published 24 January 2022)

The Fermi acceleration model describes how cosmic ray particles accelerate to great speeds by interacting with moving magnetic fields. We identify a variation of the model where light ions interact with a moving wall while undergoing pitch angle scattering through Coulomb collisions due to the presence of a heavier ionic species. The collisions introduce a stochastic component which adds complexity to the particle acceleration profile and sets it apart from collisionless Fermi acceleration models. The unusual effect captured by this simplified variation of Fermi acceleration is the nonconservation of phase space, with the possibility for a distribution of particles initially monotonically decreasing in energy to exhibit an energy peak upon compression. A peaked energy distribution might have interesting applications, such as to optimize fusion reactivity or to characterize astrophysical phenomena that exhibit nonthermal features.

DOI: [10.1103/PhysRevE.105.015207](https://doi.org/10.1103/PhysRevE.105.015207)**I. INTRODUCTION**

The Fermi acceleration model was introduced to describe how cosmic ray particles are accelerated to great speeds by interacting with moving magnetic fields [1,2]. Since then, many variations of the model have been studied. One well-known example is the Fermi-Ulam model which describes the acceleration of an ensemble of noninteracting particles bouncing between a moving wall and a stationary wall [3,4]. Several studies have examined different billiard shapes [5–8], wall movement setups [5,9–12], and particle forces [12–14] and how this affects particle acceleration and accessible points in phase space.

Consider another variation where particles interact with a moving wall while also undergoing pitch angle scattering. Suppose that the pitch angle scattering, also called Lorentz scattering, is effected by means of Coulomb collisions of light ions with a background of heavy ions. We assume that other types of collisions occur on a much greater timescale and therefore do not consider their effects in our model. We also assume that the moving wall has a negligible effect on the density of the heavy ions, either by allowing these particles to pass through or stick to the wall as it compresses. Figure 1 shows a graphic of this system. This setup is not unique in that it considers Fermi acceleration with pitch angle scattering, as other studies have investigated aspects of this collisional effect [15–18]. However, these studies have added many features simultaneously, such as electromagnetic fields, fluid effects, and complex scattering systems which do not isolate the effects we report in this paper.

The unique aspect of our problem is the simplicity of our system, paired with the limit in which we study the collisions, which results in a distinctive scaling between the change in energy and the initial energy of a particle. Since we are setting the Coulomb collision time as the smallest timescale in the system, pitch angle scattering will be the primary mechanism for reflecting particles back towards the moving wall rather

than being reflected by any wall on the other side. Introducing this stochastic effect into the system will influence the frequency at which particles interact with the moving wall and therefore also affect the total evolution of the particle distribution function. In particular, due to the relationship between the mean free path for Coulomb collisions and the speed of a particle, the rate at which a particle is accelerated by the moving wall may be heavily dependent on its initial speed and the collision frequency with the background species. This would imply that such a system could be tuned with these parameters to accelerate distributions of particles in a desired way to achieve a peaked energy distribution.

To get an idea of the underlying physics in the system, first we study a simpler one-dimensional (1D) problem where we treat Lorentz scattering by having particles reflected back towards the moving wall after traveling one collisional mean free path. We call this the plasma wall approximation, since particles are being reflected at a fixed length. This setup results in an invariant conserved during compression which significantly differs from the collisionless adiabatic invariant. In a more sophisticated model we use a random walk representation of Lorentz scattering and calculate the expected increase in energy from the wall movement. This model predicts an inverse relationship between the change in energy and the initial energy. Specifically, it predicts that in the limit of small compression ($\Delta E \ll E_0$) with many pitch angle scattering collisions ($\tau_{\text{col}} \ll T_{\text{comp}}$), the change in energy scales with $E_0^{-1/4}$, where E_0 is a particle's initial energy, τ_{col} is the time between collisions, and T_{comp} is the total compression time. We also further confirm the scaling in this limit through a basic computational particle simulation. The inverse relationship allows less energetic particles to experience a greater increase in energy than more energetic ones, resulting in narrower distributions compressed in velocity space. This is the opposite relationship of that described by the well-known example of slowly compressing a container of noninteracting




FIG. 1. Particles interacting with a wall moving at speed v_w while pitch angle scattering with a background species.

particles. This unique relationship and the non-Hamiltonian nature of the system makes this problem interesting to study, particularly because of the possibility of nonthermal features and phase space nonconservation.

The paper is organized as follows. In Sec. II we consider a simple 1D problem where particles are reflected after traveling a mean free path and identify an invariant. In Sec. III we describe the implications of our 1D random walk model and determine how the energy increase scales with a particle's initial energy. Section IV features a summary of our study and a discussion of the results.

II. 1D PLASMA WALL APPROXIMATION

Many Fermi acceleration models can be simply described by interactions of bouncing balls with moving and stationary rigid walls. Our variation is characterized by the inclusion of two species of ions with significantly different masses and the interactions between them. To get an idea of the physics of this system, we start by studying a simpler problem where particles are reflected after traveling a mean free path. This simple plasma wall approximation captures the key physics phenomena.

A. Model and assumptions

Consider a 1D model of an ensemble of ions in a box interacting with a rigid wall moving at constant speed v_w , much smaller than any particle speed. A second ensemble of more massive ions is assumed to be nearly stationary in the background and either passes through or sticks to the moving wall as it compresses. The interspecies Coulomb collision time is assumed to be the smallest timescale, followed by the total compression time and then the collision time for the light ions interacting with themselves. This time ordering prevents the light species from thermalizing during compression. In the case where there are no collisions, particles bounce back and forth between the moving wall and a stationary wall at the other end of the box, separated by distance L while conserving action. In the case where the collisional mean free path is very small compared to the system size, pitch angle scattering is considered to be the primary mechanism of reflecting particles back towards the moving wall. A simple way to model this effect in the 1D problem is to have particles be reflected after traveling one mean free path, $\lambda_{mfp} = \alpha v^4$, into the box, where




FIG. 2. Particles interacting with a moving wall while being reflected after traveling a mean free path.

α is a constant and v is the particle speed. The length of the box L is much greater than any particle's mean free path, so most particles are reflected before ever reaching the stationary wall on the other side. Therefore that wall can be neglected. Figure 2 shows a graphic of the plasma wall approximation.

B. Invariants of motion

Key aspects of the pitch angle scattering can be expressed by the 1D plasma wall model. In particular, the nature of the interaction is non-Hamiltonian, allowing for phase space nonconservation, although the system still exhibits invariants. The adiabatic invariant for collisionless particles being compressed in a 1D box with width L is the well-known

$$J_1 = vL, \quad (1)$$

and it turns out that the plasma wall model holds the invariant

$$J_2 = \Delta L + \frac{1}{4}\alpha v^4, \quad (2)$$

where ΔL is the total distance compressed, taken to be negative. The moving wall displacement over a particle bounce time is given by

$$dL = -v_w t_{\text{bounce}}, \quad (3)$$

where $t_{\text{bounce}} = 2\alpha v^3$ is the amount of time a particle with speed v spends between interactions with the moving wall. By making this substitution and using the relation $dv = 2v_w$, the equation becomes

$$dL = -\alpha v^3 dv. \quad (4)$$

Finally, by integrating and rearranging we are left with

$$\Delta L + \frac{1}{4}\alpha v^4 = C, \quad (5)$$

where C is some constant which we will denote as J_2 . Although this invariant is a direct result of v_w being a small parameter, it is not formally an adiabatic invariant. Adiabatic invariants are synonymous with action conservation in slowly varying Hamiltonian systems; however, this system is non-Hamiltonian, and we will show it does not conserve total phase space.

These two invariants have a stark physical difference, since particles which conserve J_1 will experience a greater

acceleration rate if they begin with a greater initial velocity, while the opposite is true for J_2 . This relationship occurs because in the plasma wall model, fast particles spend a greater amount of time away from the wall since the mean free path scales with v^4 , which is a phenomena unique to Lorentz scattering. The fact that less energetic particles achieve a greater increase in energy over some compression time causes velocity-space compression.

To gain insight on how an ensemble of particles evolves as a whole, we can derive a Fokker-Planck equation for the particle distribution function $f(x, v, t)$. We are most interested in how the distribution is accelerated and not necessarily in the spatial location of particles, since they will be confined within a collisional mean free path of the moving wall. Therefore the focus will instead be on deriving an equation for the spatially averaged distribution function $f(v, t)$. To further simplify the problem, we will assume that particles receive a continuous acceleration over the period of a bounce rather than a discrete gain in speed at the end of each period. This approximation is realized under the assumption that the wall speed v_w is much smaller than any particle speed and that its evolution is slow compared to a bounce time. In addition, we are only interested in the evolution of $f(v, t)$ on timescales which are much larger than the bounce time so that many bounces occur. The speed of particles in this continuous acceleration approximation will evolve according to the following stochastic Langevin equation:

$$dv = a(v)dt + \sqrt{2D(v,t)}dW, \quad (6)$$

where $a(v) = v_w/\alpha v^3$, $D(v, t)$ is a characteristic diffusion coefficient, and dW is the differential Wiener process. The Wiener process is a continuous time stochastic process which is commonly used in modeling Brownian motion [19]. In our system we have not described any physical mechanisms for diffusion; however, the inclusion of variance in the wall speed or bounce time could result in a nonzero diffusion coefficient. The Langevin equation directly implies an associated forward Kolmogorov equation for the probability distribution function $f(v, t)$, also called a Fokker-Planck equation [20]. The Fokker-Planck equation implied by the Langevin equation in (6) is

$$\frac{\partial f(v, t)}{\partial t} = -\frac{\partial}{\partial v}[a(v)f(v, t)] + \frac{\partial^2}{\partial v^2}[D(v, t)p(v, t)]. \quad (7)$$

If each particle receives a speed increase of exactly $\Delta v = 2v_w$ over the bounce time $t_{\text{bounce}} = 2\alpha v^3$ as in our model, then $D(v, t) = 0$. In this case the equation is easily solved using the method of characteristics. Solutions to this equation have the form

$$f(v, t) = v^3 h\left(v^4 - 4\frac{v_w}{\alpha}t\right), \quad (8)$$

where $h(x)$ is an arbitrary differentiable function, chosen to fit the initial condition $f_0(v, 0)$. For an initial uniform distribution between v_1 and v_2 , the solution is

$$f(v, \Delta L) = \frac{v^3 \theta\left(v^4 + 4\frac{\Delta L}{\alpha} - v_1^4\right)}{\left(v^4 + 4\frac{\Delta L}{\alpha}\right)^{3/4}} \times \left[1 - \theta\left(v^4 + 4\frac{\Delta L}{\alpha} - v_2^4\right)\right], \quad (9)$$




FIG. 3. Evolution of a step velocity distribution conserving the invariant $J_2 = \Delta L + \frac{1}{4}\alpha v^4$.

where we have made the change of variables $\Delta L = -v_w t$, and $\theta(x)$ is the usual heaviside step function. Figure 3 shows the evolution of this distribution at different stages of compression. Note from the Langevin equation that particles obey the invariant J_2 in the case of no diffusive processes during compression. Since the particles with greater speed experience less acceleration, the distribution becomes compressed in velocity space, resulting in a narrower peaked distribution.

C. Phase space volume

Although we have identified compression in velocity space, the total phase space is not necessarily compressed. As particles gain energy from interacting with the moving wall, they will occupy a greater amount of physical space since the mean free path increases with an increased speed.

To understand how these effects compete, consider an initial distribution uniformly distributed in 2D (x, v) phase space between $(v = v_0, v = v_1)$ and $(x = 0, x = \lambda_{mfp})$. When the rigid wall begins to move, particles will gain energy from interacting with this wall. As previously shown, the less energetic particles will experience the greatest increase in speed ($\Delta v_0 > \Delta v_1$). The initial and final phase space volumes are given by

$$P_i = \int_{v_0}^{v_1} \alpha v^4 dv \quad (10)$$

and

$$P_f = \int_{v_0+\Delta v_0}^{v_1+\Delta v_1} \alpha v^4 dv, \quad (11)$$

respectively. In the limit of small wall compression ($\Delta v/v \ll 1$) the change in phase space volume to first order in $\Delta v_{0,1}$ is

$$\Delta P = \alpha(v_1^4 \Delta v_1 - v_0^4 \Delta v_0). \quad (12)$$




FIG. 4. Particles interacting with a moving wall while taking a 1D random walk with a step size equal to a collisional mean free path.

In this limit we can also approximate the form of Δv as

$$\Delta v = 2v_w \frac{T_{\text{comp}}}{2\alpha v^3}. \quad (13)$$

Making this substitution into (12) yields

$$\Delta P = v_w T_{\text{comp}} (v_1 - v_0), \quad (14)$$

which clearly is greater than zero. Therefore, the expansion in physical space causes a net phase space volume increase despite the compression in velocity space.

III. RANDOM WALK MODEL

The plasma wall model described in Sec. II of ions reflecting after traveling one mean free path provides insight into the physics of Fermi acceleration with Lorentz scattering; however, it fails to fully capture the effect and cannot easily be scaled to three dimensions. A more accurate model of pitch angle scattering can be described with a random walk rather than forcing particle reflection at a mean free path. This is also relatively straightforward to extend from 1D to higher dimensions. The random walk model turns out to retain the most important feature, which is the inverse relationship between a particle's change in energy and its initial energy. Furthermore, in the limit of many collisions during the compression time, we can estimate the exact form of this relationship.

A. Additional assumptions

Particles now take a random walk with the step size equal to one mean free path rather than being reflected. After each particle collision time $\tau_{\text{col}} = \alpha v^3$, a particle will either continue on in the same direction or be reflected, exhibiting a random walk. This permits a stochastic component into the system, since a particle bounce time is now described by a probability distribution rather than a set time. A graphic of the random walk model is shown in Fig. 4.

B. 1D energy scaling

In a 1D random walk of n steps, the expected number of equalizations (or returns to the origin) r in the limit of large n




FIG. 5. Evolution of a step velocity distribution from compression while following the rate of the expected number of wall interactions given by (16).

is interestingly given by

$$E[r] = \sqrt{\frac{2}{\pi}} \sqrt{n} \quad (15)$$

[19,21]. The number of steps in our particle system can be expressed in terms of the total compression time T_{comp} and the collision time, τ_{col} , as $n = T_{\text{comp}}/\tau_{\text{col}}$. Therefore, for small compression ($\Delta v \ll v_0$) the expected increase in velocity for a particle with initial speed v_0 is

$$E[\Delta v] = \frac{2\sqrt{2}}{\sqrt{\pi}} \sqrt{\frac{T_{\text{comp}}}{\alpha v_0^3}} v_w. \quad (16)$$

This yields an unusual dependence of the speed increase with $v^{-3/2}$ or equivalently, the energy increase with $E^{-1/4}$. The inverse dependence implies that slower, less energetic particles will receive a greater kick in energy than more energetic particles, resulting in the expected velocity-space compression. Figure 5 shows the evolution of a uniform velocity distribution adhering to the expected velocity gain given by (16). Here we define $T_{\text{comp}} = -\Delta L/v_w$.

The compression of the distribution in velocity space is similar to that demonstrated in Fig. 3, although to a lesser degree. This is a result of the random walk model's weaker inverse scaling between a particle's speed and the acceleration it will experience. In this figure we have also assumed that all particles evolve according to the expected energy gain and have not accounted for the full distribution of energy gains from the random walk model. Nevertheless, we still see the expected compression in velocity space. As the distribution evolves, it becomes narrower and develops a peak in the lower energy regime of the distribution. Since the system is stochastic, the total phase space is consequently also not conserved as in the plasma wall approximation.

C. Higher dimensions

For systems of higher dimension, the scaling law given by (15) for the 1D problem is expected to hold, with the only




FIG. 6. Velocity increase following compression as a function of initial velocity. In this simulation, $\alpha = 0.1$, $v_w = 10^{-9}v_0$, and $T_{\text{comp}} = 5000\tau_{\text{col}v_0}$.

difference being the constant factor. To demonstrate this, consider an m -dimensional spatial system with the wall moving parallel to any one of the dimensions. Suppose that particles randomly walk along a uniform m -dimensional lattice with grid points separated by a mean free path distance. If each scattering direction is equally likely to occur at a step and a particle takes a total number of n steps, then the total number of these steps expected to be along the direction parallel to the wall's movement is n/m . Since the system has translation symmetry in all directions perpendicular to the wall's movement, this setup is equivalent to a 1D random walk of n/m steps, and therefore the expected number of particle-wall interactions is given by

$$E[r] = \sqrt{\frac{2}{\pi m}} \sqrt{n}. \quad (17)$$

Understandably, this is less than the strictly 1D case, since particles can now “waste” steps on other degrees of freedom. In a realistic three-dimensional (3D) system, particles are not restricted to a grid so the constant factor in (17) will be different to account for scattering at any spherical angle.

D. Numerical results

A simple particle simulation was written to investigate the effect of Coulomb pitch angle scattering in the 3D random walk case. A total of 10^5 particles were initialized at the surface of a moving, rigid wall with velocities directed away from the wall. The particles were divided into ten groups with different initial velocities in order to determine how the speed increase scaled with the initial speed. There was no stationary wall implemented on the other side of the simulation domain, so pitch angle scattering was the only mechanism responsible for turning particles back towards the moving wall. Collisions were simulated by randomly changing a particle's pitch angle every time it traveled one mean free path, $\lambda_{mfp} = \alpha v^4$. Figure 6 shows the average change in velocity for each sub-

group as a function of their initial velocity after a compression time of about 5000 collisions for the least energetic group of particles.

Clearly the result shows an inverse relationship between the two variables as shown earlier in the 1D case. When there is a significant velocity increase over the compression time, the scaling is near $\Delta v \sim v_0^{-3/2}$ (or equivalently, $\Delta E \sim E_0^{-1/4}$), as shown by the fitting curve, which is also consistent with our analytic prediction from earlier in the section. This inverse scaling is the key to obtaining the nonthermal peaked distributions shown in Figs. 3 and 5 from velocity-space compression.

IV. SUMMARY AND DISCUSSION

We presented a simple model for a two species ion ensemble interacting with a moving wall while also undergoing pitch angle scattering. We predicted an interesting inverse relationship between the change in energy from compression and the initial particle energy. It follows that less energetic particles experience greater acceleration, resulting in compression of the particle distribution in velocity space. This velocity-space compression could generate potentially favorable peaked energy distributions as shown in Figs. 3 and 5.

The nonthermal phenomena predicted by these models could be of interest to various areas of plasma physics, since we are considering Lorentz scattering of charged particles. In particular, due to the mass difference between the two species, the model could describe some aspects of $p^{11}\text{B}$ interactions. This is a potential fuel source for aneutronic fusion [22], and the velocity-space compression could provide a mechanism for obtaining favorable proton energy distributions to increase fusion reactivity [23–25]. The moving wall in this case could represent either a physical wall compressing or a moving magnetic field structure in a magnetic mirror confinement setup [26]. The model could also be used to describe a variation of cosmic ray acceleration, where the Lorentz scattering mean free path is small compared to the system size. Identifying acceleration mechanisms in astrophysical settings using Fermi acceleration models is an ongoing area of study [27–30], and the phenomena outlined in this paper may be applicable to the field.

We opted for simplicity in our models to isolate the important effects described in this paper. However, in real plasmas there are more complex features such as collisions within a species which allows ions to thermalize. Thermalization could significantly dampen velocity space compression, which is why we examined the limit where interspecies collisions dominate over the light ion collisions with themselves. In this limit we have identified some unique effects of Fermi acceleration with Lorentz scattering. The most interesting aspect of the results is the potential for nonthermal, non-Hamiltonian features in the compression due to the nonconservation of phase space.

ACKNOWLEDGMENTS

The authors are thankful to E. Kolmes, I. Ochs, and T. Rubin for helpful conversations. This work is supported by NSF PHY-1805316 and NNSA DE-SC0021248.

- [1] E. Fermi, *Phys. Rev.* **75**, 1169 (1949).
- [2] M. Lemoine, *Phys. Rev. D* **99**, 083006 (2019).
- [3] S. Ulam, *Proceedings of the 4th Berkeley Symposium on Mathematical Statistics and Probability* (California University Press, Berkeley, CA, 1961), p. 315
- [4] A. Lichtenberg and M. Lieberman, *Regular and Chaotic Dynamics*, 2nd ed. (Springer, New York, 1992).
- [5] V. Gelfreich, V. Rom-Kedar, and D. Turaev, *Chaos* **22**, 033116 (2012).
- [6] J. Zhou, *Nonlinearity* **33**, 1542 (2020).
- [7] F. Lenz, F. K. Diakonov, and P. Schmelcher, *Phys. Rev. Lett.* **100**, 014103 (2008).
- [8] E. D. Leonel, D. F. M. Oliveira, and A. Loskutov, *Chaos* **19**, 033142 (2009).
- [9] C. Jarzynski, *Phys. Rev. E* **48**, 4340 (1993).
- [10] A. Loskutov, A. B. Ryabov, and L. G. Akinshin, *J. Phys. A: Math. Gen.* **33**, 7973 (2000).
- [11] V. Gelfreich and D. Turaev, *J. Phys. A: Math. Theor.* **41**, 212003 (2008).
- [12] C. K. Langer and B. N. Miller, *Chaos* **25**, 073114 (2015).
- [13] E. D. Leonel and P. V. E. McClintock, *J. Phys. A: Math. Gen.* **38**, 823 (2005).
- [14] Felipe Augusto O. Silveira, S. G. Alves, E. D. Leonel, and D. G. Ladeira, *Phys. Rev. E* **103**, 062205 (2021).
- [15] B. D. G. Chandran, *Phys. Rev. Lett.* **85**, 4656 (2000).
- [16] R. Selkowitz and E. G. Blackman, *MNRAS* **354**, 870 (2004).
- [17] T. Liu III, S. Lu, V. Angelopoulos, H. Hietala, and L. B. Wilson III, *J. Geophys. Res. Space Phys.* **122**, 9248 (2017).
- [18] J. Scott, W. Cocke, R. Chevalier, and D. Wentzel, *Astrophys. Space Sci.* **53**, 421 (1978).
- [19] W. Feller, *An Introduction to Probability Theory and Its Applications* (Wiley, New York, 1950), Vol. 1.
- [20] H. Risken and H. Haken, *The Fokker-Planck Equation: Methods of Solution and Applications*, 2nd ed. (Springer, New York, 1989).
- [21] C. Grinstead and J. L. Snell, *Introduction to Probability* (American Mathematical Society, Providence, RI, 1997).
- [22] J. Dawson, in *Fusion*, edited by E. Teller (Academic Press, New York, 1981), Vol. 1, pp. 453–501.
- [23] H. W. Becker, C. Rolfs, and H. P. Trautvetter, *Z. Phys. A* **327**, 341 (1987).
- [24] W. M. Nevins and R. Swain, *Nucl. Fusion* **40**, 865 (2000).
- [25] M. H. Sikora and H. R. Weller, *J. Fusion Energy* **35**, 538 (2016).
- [26] W. M. Nevins, *J. Fusion Energy* **17**, 25 (1998).
- [27] M. A. Malkov and L. O. Drury, *Rep. Prog. Phys.* **64**, 429 (2001).
- [28] W. Fox, J. Park, W. Deng, G. Fiksel, A. Spitkovsky, and A. Bhattacharjee, *Phys. Plasmas* **24**, 092901 (2017).
- [29] A. Veltri and V. Carbone, *Phys. Rev. Lett.* **92**, 143901 (2004).
- [30] V. Petrosian and A. Bykov, *Space Sci. Rev.* **134**, 207 (2008).