

Multi-volume, partially-relaxed MHD equilibria

S.R. Hudson, R.L. Dewar^a, M.J. Hole^a, M. McGann^a

Princeton Plasma Physics Laboratory, P.O. Box 451, Princeton, NJ, 08543, USA

^aPlasma Research Laboratory, Research School of Physics and Engineering, The Australian National University, Canberra ACT 0200, Australia

shudson@pppl.gov

The commonly used equation of ideal force balance, $\nabla p = \mathbf{j} \times \mathbf{B}$, is pathological when the magnetic field, \mathbf{B} , is chaotic. This is because $\mathbf{B} \cdot \nabla p = 0$ implies that any continuous pressure, p , will be a devils staircase without a well defined gradient. This pathological structure causes problems for the so-called Spitzer iterative approach, which is fundamentally ill-posed as it depends on inverting magnetic differential equations and such equations have a dense set of singularities. We suggest instead a well-posed equilibrium construction based on a constrained energy principle.

Consider [1] a plasma region comprised of a set of N nested annular regions which are separated by a discrete set of toroidal interfaces, \mathcal{I}_l . In each volume, \mathcal{V}_l , bounded by the \mathcal{I}_{l-1} and \mathcal{I}_l interfaces, the plasma energy, U_l , the global-helicity, H_l , and the “mass”, M_l , are given by the integrals:

$$U_l = \int_{\mathcal{V}_l} \left(\frac{p}{\gamma - 1} + \frac{B^2}{2\mu_0} \right) dv, \quad H_l = \int_{\mathcal{V}_l} \mathbf{A} \cdot \mathbf{B} dv, \quad M_l = \int_{\mathcal{V}_l} p^{1/\gamma} dv, \quad (1)$$

where \mathbf{A} is the vector potential, $\mathbf{B} = \nabla \times \mathbf{A}$.

The equilibrium states that we seek minimize the total plasma energy, subject to the constraints of conserved helicity and conserved mass in each annular region. We allow arbitrary variations in both the magnetic field in each annulus and the geometry of the interfaces, except that we assume the magnetic field remains tangential to the interfaces which we consider to act as ideal barriers.

The Euler-Lagrange equations show [1] that that in each annulus the magnetic field satisfies $\nabla \times \mathbf{B} = \mu \mathbf{B}$, and across each interface the total pressure is continuous, $[[p + B^2/2]] = 0$.

We have implemented this model in a code, the Stepped Pressure Equilibrium Code (SPEC), and numerical results will be presented. The SPEC code uses a mixed Fourier-Finite element representation for the vector potential. Quintic polynomial basis functions give rapid convergence in the radial discretization, and the poloidal angle is adjusted to minimize a “spectral-width”. For given interface geometries the Beltrami fields in each annulus are constructed in parallel, and a Newton method (with quadratic-convergence) is implemented to adjust the interface geometry to satisfy force-balance.

[1] Relaxed plasma equilibria and entropy-related plasma self-organization principles, Dewar *et al.*, Entropy 10:621, 2008.